

Spherical conjugacy classes and involutions in the Weyl group

Giovanna Carnovale

Dipartimento di Matematica Pura ed Applicata
Torre Archimede - via Trieste 63 - 35121 Padova - Italy
email: carnoval@math.unipd.it

Abstract

Let G be a simple algebraic group over an algebraically closed field of characteristic zero or positive odd, good characteristic. Let B be a Borel subgroup of G . We show that the spherical conjugacy classes of G intersect only the double cosets of B in G corresponding to involutions in the Weyl group of G . This result is used in order to prove that for a spherical conjugacy class $\mathcal{O}$ with dense B -orbit $v_0 \subset BwB$ there holds $\ell(w) + \text{rk}(1 - w) = \dim \mathcal{O}$ extending a characterization of spherical conjugacy classes obtained by Cantarini, Costantini and the author to the case of groups over fields of odd, good characteristic.

Introduction

If an algebraic group acts with finitely many orbits, a natural way to understand the action is given by the combinatorics of the Zariski closures of such orbits. In [19], [23], a detailed description of the combinatorics of the closures of orbits for a Borel subgroup B in a symmetric space G/K is given. The description is provided in terms of an action, on the set of these orbits, of a monoid $M(W)$ related to the Weyl group W of G . This action is best understood considering the decomposition into B -orbits of an orbit of a minimal parabolic subgroup. Through this approach several invariants of the B -orbits can be determined, including their dimension. To each B -orbit it is possible to associate a Weyl group element and the Weyl group element corresponding to the (unique) dense B -orbit in the symmetric space can be described in combinatorial terms. A formula for the dimension of each B -orbit is provided in terms of its associated Weyl group element and the sequence of elements in the monoid that are necessary to reach it from a closed B -orbit. When the symmetric space corresponds to an inner involution, that is, if it corresponds

to a conjugacy class in G , the attached Weyl group element is just the element corresponding to the Bruhat cell containing the B -orbit.

The monoid action can be carried over to homogeneous spaces of algebraic groups for which the action of the Borel subgroup has finitely many orbits, i.e., the spherical homogeneous spaces ([22]) and it can be used to define representations of the Hecke algebra ([15]). A more geometric approach to a Bruhat order on spherical varieties has been addressed in [6]. Besides, a genuine Weyl group action on the set of B -orbits on a spherical homogeneous space was defined in [14].

The action of $M(W)$ on a spherical homogeneous space does not afford all nice properties that it had in the symmetric case (see [7] for a few key counterexamples) and it is natural to ask which properties still hold for spherical conjugacy classes. One of the main differences between the general spherical case and the symmetric case is that there are B -orbits that do not lie in the orbit of a closed one when acting by $M(W)$. However, every B -orbit can be reached from a closed B -orbit through a sequence of moves involving either the $M(W)$ -action or the W -action ([22]).

A natural question is whether we can provide formulas for the dimension of each B -orbit in a spherical conjugacy class in terms of the actions of $M(W)$ and W . Although not all results in [19] hold at this level of generality, there are properties that hold true in general. For instance, the dimension of the dense B -orbit in a spherical conjugacy class is governed by a formula analogous to the formula for the dimension of a B -orbit in a symmetric conjugacy class. This result was achieved in [8], leading to a characterization of spherical conjugacy classes in simple algebraic groups. The interest in this formula lied in the verification of De Concini-Kac-Procesi conjecture on the dimension of irreducible representations of quantum groups at the roots of unity ([9]) in the case of spherical conjugacy classes. For this reason, the analysis was restricted to the case of an algebraic group over an algebraically closed field of characteristic zero. In order to obtain the characterization, a classification of all spherical conjugacy classes in a simple algebraic group was needed, and part of the results were obtained through a case-by-case analysis involving this classification.

In the present paper we apply the combinatorics of B -orbit closures for spherical conjugacy classes in simple algebraic groups to retrieve the formula in [8]. This will show that the characterization of spherical conjugacy classes can be achieved without making use of their classification and without drastic restrictions on the characteristic of the base field.

A first question to be answered concerns which Weyl group elements may correspond to a B -orbit in a spherical conjugacy class. In the case of a symmetric conjugacy class it is immediate to see that such Weyl group elements are involutions. An analysis of the actions of $M(W)$ and W allows us to generalize this result to all spherical conjugacy classes.

Theorem 1 *All B -orbits in a spherical conjugacy class lie in Bruhat cells corresponding to involutions in the Weyl group.*

In order to understand the Weyl group elements associated with the dense B -orbit we analyze the variation of the Weyl group element with respect to the action of the monoid $M(W)$. This analysis leads to a description of the stationary points, i.e., of those B -orbits for which the associated Weyl group element does not change under the action of all standard generators of $M(W)$. Stationary point different from the dense B -orbit cannot exist in symmetric conjugacy classes but they exist, for instance, in spherical unipotent conjugacy classes.

The results in [23] allow us to describe the Weyl group element corresponding to a stationary point, and more precisely, the one associated with the dense B -orbit.

Combining the analysis of representatives of the dense B -orbit with results in [7] yields a new proof of the formula in [8], that holds now in almost all characteristics and does not require the classification of spherical conjugacy classes:

Theorem 2 *Let $\mathcal{O}$ be a spherical conjugacy class in a simple algebraic group G , let v_0 be its dense B -orbit and let BwB be the Bruhat double coset containing v_0 . Then $\dim \mathcal{O} = \ell(w) + \text{rk}(1 - w)$.*

It is proved in [8] with a characteristic-free argument that if a conjugacy class $\mathcal{O}$ intersects some BwB with $\ell(w) + \text{rk}(1 - w) = \dim \mathcal{O}$ then $\mathcal{O}$ is spherical, hence the results in the present paper provide a characteristic-free proof of the characterization of spherical conjugacy class given in [8].

Acknowledgements

This research was partially supported by Marie Curie Research Training Network LIEGRITS Flags, Quivers and Invariant Theory in Lie Representation Theory number MRTN-CT 2003-505078.

1 Preliminaries

Let G be a simple algebraic group over an algebraically closed field k of characteristic 0 or odd and good ([24, §4.3]). Let B be a Borel subgroup of G , T a maximal torus contained in B and B^- the Borel subgroup opposite to B . Let U (respectively U^-) be the unipotent radical of B (respectively B^-). For a group K we shall denote by K° the connected component of the identity of K .

We shall denote by Φ the set of roots relative to (B, T) , by Φ^+ the corresponding positive roots, by $\Delta = \{\alpha_1, \dots, \alpha_n\}$ the corresponding set of simple roots. We shall use the numbering of the simple roots in [3, Planches I-IX]. The usual inner

product between the roots α and β will be denoted by (α, β) . If the root system is simply laced we will consider all its roots as long roots. We shall indicate by P^+ and Q , respectively, the set of dominant weights and the root lattice associated with Φ . An element of T corresponding to the co-root $\alpha^\vee$ will be indicated by $\alpha^\vee(h)$.

By Π we shall always denote a subset of Δ and $\Phi(\Pi)$ will indicate the corresponding root subsystem.

We shall denote by W be the Weyl group associated with G and by s_α the reflection corresponding to the root α . By $\ell(w)$ we shall denote the length of the element $w \in W$ and by $\text{rk}(1 - w)$ we shall mean the rank of $1 - w$ in the standard representation of the Weyl group. By w_0 we shall denote the longest element in W and by ϑ we shall denote the automorphism of Φ given by $-w_0$. If $\Pi \subset \Delta$ we shall denote by W_Π the parabolic subgroup of W generated by the simple reflections in Π . If $N(T)$ is the normalizer of T in G then $W = N(T)/T$; given an element $w \in W$ we shall denote a representative of w in $N(T)$ by $\dot{w}$. For any root α of Φ we shall denote by $x_\alpha(t)$ the elements of the corresponding root subgroup X_α of G . We shall choose the representatives $\dot{s}_\alpha \in N$ of the reflection $s_\alpha \in W$ as in [21, Theorem 8.1.4].

We assume that we have fixed an ordering of the positive roots so that every $u \in U$ is written uniquely as an ordered product of element of the form $x_\alpha(t)$, for $t \in k$ and $\alpha \in \Phi^+$. Given an element $u \in U$ (respectively U^-), by abuse of language we will say that a root $\gamma \in \Phi^+$ (respectively $-\Phi^+$) *occurs in u* if for the expression of u as an ordered product of $x_\alpha(t_\alpha)$'s we have $t_\gamma \neq 0$.

If $\alpha \in \Delta$ we shall indicate by P_α the minimal non solvable parabolic subgroup containing $X_{-\alpha}$ and by P_α^u its unipotent radical.

For $w \in W$, we shall denote by U^w (respectively, U_w) the subgroup generated by the root subgroups X_α corresponding to those $\alpha \in \Phi^+$ for which $w^{-1}(\alpha) \in -\Phi^+$ (respectively, Φ^+). We shall denote by T^w the subgroup of the torus that is centralized by any representative $\dot{w}$ of w .

Given an element $x \in G$ we shall denote by $\mathcal{O}_x$ the conjugacy class of x in G and by G_x (resp. B_x , resp. T_x) the centralizer of x in G (resp. B , resp. T). For a homogeneous space G/H we shall denote by $\mathcal{V}$ the set of B -orbits in G/H .

Definition 1.1 *Let K be a connected algebraic group over k and let H be a closed subgroup of K . The homogeneous space K/H is called spherical if there exists a Borel subgroup of K with a dense orbit.*

It is well-known ([5], [25] in characteristic 0, [10], [14] in positive characteristic) that, if G/H is a spherical homogeneous space the set $\mathcal{V}$ of B -orbits in G/H is finite.

2 B -orbits and Bruhat decomposition

Let $\mathcal{O}$ be a conjugacy class of G and let $\mathcal{V}$ be the set of B -orbits in $\mathcal{O}$. There is a natural map $\phi: \mathcal{V} \rightarrow W$ associating to $v \in \mathcal{V}$ the element w in the Weyl group of G for which $v \subset BwB$. The set $\mathcal{V}$ carries a partial order given by: $v \leq v'$ if $\overline{v} \subset \overline{v'}$. When $\mathcal{O}$ is spherical the minimal B -orbits are the closed ones and there is a unique maximal orbit, namely the dense B -orbit v_0 in $\mathcal{O}$.

Lemma 2.1 *Let $\mathcal{O}$ be a conjugacy class and let $v, v' \in \mathcal{V}$. If $v \leq v'$ then $\phi(v) \leq \phi(v')$ in the Bruhat order in W .*

Proof. We have: $v \subset \overline{v} \subset \overline{v'} \subset \overline{B\phi(v')B} = \dot{\cup}_{\sigma \leq \phi(v')} B\sigma B$ so $\phi(v) \leq \phi(v')$. $\square$

Lemma 2.2 *Let $x \in G$ be either semisimple or unipotent and let $\mathcal{O}_x$ be a spherical conjugacy class. The image through ϕ of a closed B -orbit in $\mathcal{O}$ is 1.*

Proof. By [22, §3.4 (b)] if $\mathcal{O} = \mathcal{O}_x$ and $H = G_x$, the closed B -orbits correspond to those BgH/H with BgH closed in G , so that $B_H = (H \cap g^{-1}Bg)^\circ$ is a Borel subgroup of H and of H° . Let x be semisimple. Since it is not restrictive to assume that G is simply connected, we have $H = H^\circ$ and $x \in Z(H) = Z(B_H)$ by [21, Corollary 6.2.9]. Hence, the representative gxg^{-1} of the closed B -orbit lies in B .

Let x be unipotent. By [24, §3.15] with $S = \{x\}$ the element $x \in H^\circ$, hence $x \in Z(H^\circ) \subset B_H$ and we have the statement. $\square$

Remark 2.3 All closed B -orbits in a spherical conjugacy class $\mathcal{O}_x$ have the same dimension ([22, §3.4 (b)]) namely $\dim B - \dim B_{G_x}$ where B_{G_x} denotes a Borel subgroup in the centralizer of x .

Remark 2.4 The converse of Lemma 2.2 does not hold for spherical unipotent elements. For instance, if $\mathcal{O}$ is a spherical unipotent conjugacy class in $G = SL_n(\mathbb{C})$ the combinatorics of the closures of the B -orbits that are contained in B is described in [16]: if $\mathcal{O}$ corresponds to the minimal nilpotent orbit in G the B -orbits that are contained in B are in bijection with the traspositions in S_n , and only one, namely $B.x_{\beta_1}(1) = X_{\beta_1} \setminus \{1\}$, is closed.

Remark 2.5 In a spherical semisimple conjugacy class, v being closed is equivalent to $\phi(B.x) = 1$ and to $v \cap T \neq \emptyset$. Indeed if v is closed then $v \subset B$ so a representative $x \in v$ is conjugate in B to some element in T . Viceversa, if $v = B.t$ for some $t \in T$ then t normalizes B and by [2, Theorem 9.2] the conjugacy class $B.t$ is closed.

Let $M = M(W)$ be the monoid with elements $m(w)$ indexed by the elements $w \in W$ with relations

$$m(s)m(w) = m(sw), \text{ if } l(sw) > l(w), \quad m(s)m(w) = m(w), \text{ if } l(sw) < l(w).$$

The monoid $M(W)$ is generated by the elements $m(s)$ corresponding to simple reflections, subject to the braid relations and to the relation $m(s)^2 = m(s)$. In [19] an action of the monoid $M(W)$ on the set of B -orbits of a symmetric space G/K is defined. This action can be generalized to an action of $M(W)$ on the set $\mathcal{V}$ of B -orbits of a spherical homogeneous space (see, for instance, [22, §3.6]). The action of $m(s)$, for a simple reflection $s = s_\alpha$ is given as follows. If P_α is the minimal parabolic subgroup corresponding to α and $v \in \mathcal{V}$ then $m(s).v$ is the dense B -orbit in $P_\alpha v$. This action is analyzed in [7], [14], [15, §4.1], [19]. We provide an account of the information we will need.

Given $v \in \mathcal{V}$, choose $y \in v$ with stabilizer $(P_\alpha)_y$ in P_α . Then $(P_\alpha)_y$ acts on $P_\alpha/B \cong \mathbb{P}^1$ with finitely many orbits. Let $\psi: (P_\alpha)_y \rightarrow PGL_2(k)$ be the corresponding group morphism. The kernel of ψ is $\text{Ker}(\alpha)P_\alpha^u$.

The image H of $(P_\alpha)_y$ in $PGL_2(k)$ is either: $PGL_2(k)$; or solvable and contains a nontrivial unipotent subgroup; or a torus; or the normalizer of a torus. Here is a list of the possibilities that may occur:

- I $P_\alpha v = v$ so $H = PGL_2(k)$;
- IIa $P_\alpha v = v \cup m(s)v$, with $\dim v = \dim P_\alpha v - 1$. In this case we may choose an $y \in v$ for which $\psi(X_\alpha) \subset H \subset \psi(B)$.
- IIb $P_\alpha v = v \cup v'$, with $\dim v' = \dim v - 1$ and v is open in $P_\alpha v$ so $m(s)v = v$. In this case we may choose an $y \in v$ for which $\psi(X_{-\alpha}) \subset H \subset \psi(B^-)$.
- IIIa $P_\alpha v = v \cup v' \cup m(s)v$, with $\dim v = \dim v' = \dim P_\alpha v - 1$ and $v \neq v'$. In this case we may choose an $y \in v$ for which $H = \psi(T)$. Then $\dot{s}_\alpha y \dot{s}_\alpha^{-1} \in v'$ and $\dot{s}_\alpha x_\alpha(t) y x_\alpha(-t) \dot{s}_\alpha^{-1} \in m(s)v$ for $t \neq 0$.
- IIIb $P_\alpha v = v \cup v' \cup v''$, with $\dim v - 1 = \dim v' = \dim v''$ and v is open in $P_\alpha v$ so $m(s)v = v$. In this case we may choose an $y \in v$ for which $H = \psi(\dot{s}_\alpha x_\alpha(-1) T x_\alpha(1) \dot{s}_\alpha^{-1})$.
- IVa $P_\alpha v = v \cup m(s)v$, with $\dim v = \dim P_\alpha v - 1$. We may choose an $y \in v$ for which $H = \psi(N(T))$. Besides, $\dot{s}_\alpha y \dot{s}_\alpha^{-1} \in v$ and $\dot{s}_\alpha x_\alpha(t) y x_\alpha(-t) \dot{s}_\alpha^{-1} \in m(s)v$ for $t \neq 0$.
- IVb $P_\alpha v = v \cup v'$, with $\dim v = \dim v' + 1$, and v is open in $P_\alpha v$ so $m(s)v = v$. In this case we may choose an $y \in v$ for which $H = \psi(N(\dot{s}_\alpha T \dot{s}_\alpha^{-1}))$.

Based on the structure of H , cases II, III, and IV are also called type U , type T and type N , respectively.

Based on this analysis a W -action on $\mathcal{V}$ can be defined ([14], [15, §4.2.5, Remark]) as follows: in case II the two B -orbits are interchanged; in case III the two non-dense orbits are interchanged, in all other cases the B -orbits are fixed. The image of $v \in \mathcal{V}$ through the action of a simple reflection $s \in W$ will be denoted by $s.v$.

We recall ([22, §3.6]) that a *reduced decomposition* of $v \in \mathcal{V}$ is a pair $(\mathbf{v}, \mathbf{s})$ with $\mathbf{v} = (v(0), v(1), \dots, v(r))$ a sequence of distinct elements in $\mathcal{V}$ and $\mathbf{s} = (s_{i_1}, \dots, s_{i_r})$ a sequence of simple reflections such that $v(0)$ is closed and $v(j) = m(s_{i_j}).(v(j-1))$ for $1 \leq j \leq r-1$.

All B -orbits in a symmetric homogeneous space admit a reduced decomposition ([19, §7]). This is still the case for the dense B -orbit in spherical homogeneous spaces but it is not always the case for generic B -orbits in a spherical homogeneous space. The reader can refer to [7] for a series of counterexamples. We will use a weaker notion of decomposition that exists for every $v \in \mathcal{V}$.

Given a reduced decomposition $(\mathbf{v}, \mathbf{s}) = ((v(0), \dots, v(r)), \mathbf{s})$ of $v \in \mathcal{V}$ a *subexpression* of $(\mathbf{v}, \mathbf{s})$ ([22, §3.6]) is a sequence $\mathbf{x} = (v'(0), v'(1), \dots, v'(r))$ of elements in $\mathcal{V}$ with $v'(0) = v(0)$ and such that for $1 \leq i \leq r$ only one of the following alternatives occurs:

- (a) $v'(j-1) = v'(j)$;
- (b) $v'(j-1) \neq v'(j)$, $\dim v'(j-1) = v'(j)$ and $v'(j) = s_j.v'(j-1)$;
- (c) $\dim v'(j-1) = \dim v'(j) - 1$ and $v'(j) = m(s_j).(v'(j-1))$.

The element $v'(r)$ is called the *final term* of the subexpression. Even though some B -orbits in a spherical homogeneous space might not have a reduced decomposition, every $v \in \mathcal{V}$ is the final term of a subexpression of a reduced decomposition of the dense B -orbit v_0 . This is to be found in [22, §3.6 Proposition 2] and it holds also in positive odd characteristic.

Lemma 2.6 *Let $\mathcal{O}$ be a spherical conjugacy class, let $v \in \mathcal{V}$ and let s be a simple reflection. If $w = \phi(v)$ is an involution then $w' = \phi(m(s).v)$ is an involution.*

Proof. We consider $P_\alpha v = v \cup (BsB).v$. If $m(s)v = v$ there is nothing to prove. Let us assume that $m(s)v \subset (BsB).v$. Then $m(s).v \subset BsBwBsB$. The following four possibilities may occur:

1. $l(sws) = l(w) + 2$: by [21, Lemma 8.3.7] we have $w' = sws$ and the statement holds.

2. $l(sw) > l(w)$ and $l(sws) = l(w)$: since w is an involution, by [23, Lemma 3.2 (ii)] for $\theta = \text{id}$ we have $sw = ws$. Besides, by [21, Lemma 8.3.7] we have

$$m(s).v \subset BswBsB = BswB \cup BswsB.$$

hence $w' \in \{sw, w\}$ is an involution.

3. $l(sw) < l(w)$ and $l(sws) = l(w)$: again by [23, Lemma 3.2 (i)] for $\theta = \text{id}$ we have $sw = ws$. Then

$$m(s).v \subset BsBwBsB \subset BswsB \cup BswB$$

so $w' \in \{w, ws\}$ is an involution.

4. $l(sws) = l(w) - 2$: we have

$$m(s).v \subset BsBwBsB \subset BswB \cup BswsB \cup BwsB \cup BwB.$$

By [22, §3.6 Proposition 1 (a)] we have $v \leq m(s).v$ so by Lemma 2.1 there holds $w \leq w'$ hence $w' \neq sw, ws, sws$ and $w' = w$ is an involution. $\square$

Corollary 2.7 *Let $\mathcal{O}_x$ be a spherical conjugacy class with x either semisimple or unipotent and let $v \in \mathcal{V}$ admit a reduced decomposition. Then $\phi(v)$ is an involution. In particular, this holds for the dense B -orbit of a spherical semisimple or of a spherical unipotent conjugacy class.*

Proof. The first assertion follows by induction on $\dim v$ using Lemma 2.2 and Lemma 2.6. The last statement follows from [22, §3.6, Proposition 3(ii)]. $\square$

Theorem 2.8 *Let $\mathcal{O}$ be a spherical conjugacy class, and let $\phi: \mathcal{V} \rightarrow W$ be the natural map. Then the image of ϕ consists of involutions.*

Proof. We first consider a spherical semisimple conjugacy class. Let $v \in \mathcal{V}$ and let x be a subexpression of a reduced decomposition of the dense B -orbit v_0 with initial term a closed B -orbit $v(0)$ and final term v . We proceed by induction on $\dim v$. If v has minimal dimension then it is closed, otherwise it would contain in its closure a B -orbit of strictly smaller dimension. It follows from Lemma 2.2 that $\phi(v) = 1$.

Let $v = v(r)$ not be closed so that $\dim v(r) > \dim v(0)$. If $v(r) = v(r-1)$ we may shorten the subexpression replacing r by $r-1$. Hence we may assume that $v(r) = m(s)v(r-1)$ or $v(r) = s.v(r-1)$ for some simple reflection s . If $v(r) = m(s)v(r-1)$ we may use Lemma 2.6. If $v(r) = s.v(r-1)$, then

$\dim(v(r-1)) = \dim v(r)$. If we proceed downwards along the terms of the subexpression we might have a sequence of steps in which either the B -orbit does not change or it changes through the W -action, but we will eventually reach a step at which $v(j) = m(s')(v(j-1))$ with $\dim(v(j)) > \dim v(j-1)$, where we can apply Lemma 2.6. Hence there is a B -orbit v' in the sequence with $\dim v' = \dim v$, and $\phi(v')$ is an involution w . Therefore we may reduce to the case in which $v' = v(r-1)$ and $v = v(r) = s.v(r-1)$. The analysis of the decompositions into B -orbits of $P_s v$ shows that we are in case IIIa. Then $v \in Bsv'sB \subset BsBwBsB$.

If $\ell(sws) = \ell(w) + 2$ then $v \in BswsB$ and we have the statement. If $sw > w$ and $\ell(sws) = \ell(w)$ then $sw = ws$ and $v \in BswB \cup BswsB$ so $\phi(v)$ is an involution. If $sw < w$ and $\ell(w) = \ell(sws)$ then $sw = ws$ and $v \in BwB \cup BswB$ so $\phi(v)$ is an involution. Let us assume that $\ell(sws) = \ell(w) - 2$ with $s = s_\alpha$. It follows from the proof of Lemma 2.1 in this case that $\phi(m(s)v) = \phi(m(s)v') = w = \phi(v')$. By [12, Lemma 1.6] we have $w\alpha \in -\Phi^+$ so we may choose representatives $x = u_x \dot{w} v_x$ and $y = u_y \dot{w} v_y$ of the same B -orbit v' with $u_x, u_y \in U^w$, $v_x, v_y \in U$ for which α does not occur in u_x and v_y .

Conjugation by $\dot{s}_\alpha$ maps x in $BswB \cup BswsB$, hence $\dot{s}_\alpha x \dot{s}_\alpha^{-1} \in v$ and $\phi(v) \in \{sw, sws\}$. On the other hand, conjugation by $\dot{s}_\alpha$ maps y in $BswsB \cup BwsB$, hence $\dot{s}_\alpha y \dot{s}_\alpha^{-1} \in v$ and $\phi(v) \in \{sw, sws\} \cap \{ws, sws\}$ is an involution. Thus we have the statement for spherical semisimple conjugacy classes.

Let us consider the spherical conjugacy class of an element $x \in G$ with Jordan decomposition su . The proof will follow by induction once we show that the image through ϕ of a closed B -orbit is an involution. As in the proof of Lemma 2.2, if $y = gxg^{-1}$ is a representative of a closed B -orbit then $(G_x \cap g^{-1}Bg)^\circ$ is a Borel subgroup of G_x° and $u \in Z(G_x^\circ)$ by [24, §3.15]. Thus the unipotent part gug^{-1} of y lies in B and $\phi(B.y) = \phi(B.gsg^{-1})$. The conjugacy class $\mathcal{O}_s$ is spherical because it can be identified with G/G_s that is a quotient of $G/G_s \cap G_u = G/G_x$. By the first part of the proof $\phi(B.y)$ is an involution. $\square$

Remark 2.9 The reader is referred to [8, §1.4, Remark 4] for a different proof, in characteristic zero, that the image through ϕ of the dense B -orbit v_0 , denoted by $z(\mathcal{O})$, is an involution. In the same paper, the image of $\phi(w_0)$ of all spherical conjugacy classes of a simple algebraic group over $\mathbb{C}$ is explicitly computed.

3 Stationary points

In this Section we shall analyze those elements in $v \in \mathcal{V}$ for which $\phi(m(s)v) = \phi(v)$ for all simple reflections $s \in W$.

Definition 3.1 Let $v \in \mathcal{V}$, let $w = \phi(v)$ and let α be a simple root. We say that

v is a stationary point with respect to α if $\phi(v) = \phi(m(s_\alpha)v)$. We say that v is a stationary point if it is a stationary point with respect to all simple roots.

Stationary points different from the dense B -orbit do not exist in symmetric conjugacy classes by the results in [19] but they do exist in unipotent spherical conjugacy classes:

Example 3.2 Let $\mathcal{O}_{\min}$ be the minimal nontrivial unipotent conjugacy class in a group G of semisimple rank at least 2. It is well-known that $\mathcal{O}_{\min}$ is spherical. If β denotes the highest root in Φ then $B.x_\beta(1) = X_\beta \setminus \{1\}$ is a stationary point. Indeed, if α is a simple root $P_\alpha = Bs_\alpha X_\alpha \cup B$ and $P_\alpha.x_\beta(t) \subset B.X_{s_\alpha(\beta)} \cup B.x_\beta(1) \subset B$ so $\phi(B.x_\beta(1)) = \phi(m(s_\alpha)(B.x_\beta(1))) = 1$.

The following lemmas describe stationary points with respect to a simple root.

Lemma 3.3 Let $v \in \mathcal{V}$ with $w = \phi(v)$. Let α be a simple root such that $s_\alpha w < w$ in the Bruhat order. Then v is a stationary point with respect to α .

Proof. Let us put $s = s_\alpha$. If $sw < w$ then $\phi(m(s)v) \in \{sws, sw, ws, w\}$ and $\phi(m(s)v) \geq \phi(v)$. If $sw = ws$ the statement follows because $ws < w$ and $sws = w$. Otherwise it follows because $sw, ws, sws < w$. $\square$

Lemma 3.4 Let $v \in \mathcal{V}$ with $w = \phi(v)$. Let α be a simple root such that $s_\alpha w > w$ in the Bruhat order. Let $x = u\dot{w}v \in v$ with $u \in U^w$, $t \in T$, and $v \in U$. Then v is a stationary point with respect to α if and only if the following conditions hold:

1. $s_\alpha w = ws_\alpha$;
2. $v \in P_\alpha^u$, the unipotent radical of P_α ;
3. $X_{\pm\alpha}$ commutes with $\dot{w} \in N(T)$.

Proof. Let $v \in \mathcal{V}$ be a stationary point with respect to $\alpha \in \Delta$, let $x \in v$ with unique decomposition $x = u\dot{w}v$ and let $s_\alpha w > w$. It follows from Theorem 2.8 that w is an involution. We have either $s_\alpha ws_\alpha > s_\alpha w$ or $s_\alpha ws_\alpha = w$. If the first case were possible, we would have $\dot{s}_\alpha x \dot{s}_\alpha^{-1} \in Bs_\alpha ws_\alpha B$ and v would not be a stationary point. Hence 1 holds, $w\alpha = \alpha$ and α does not occur in $u \in U^w$.

Let us consider $y = \dot{s}_\alpha x \dot{s}_\alpha^{-1}$. The element

$$y = (\dot{s}_\alpha u \dot{s}_\alpha^{-1})(\dot{s}_\alpha \dot{w} \dot{s}_\alpha^{-1})(\dot{s}_\alpha v \dot{s}_\alpha^{-1}) \in Bw(\dot{s}_\alpha v \dot{s}_\alpha^{-1}).$$

If α would occur in v then by [21, Lemmas 8.1.4, 8.3.7] we would have $y \in Bws_\alpha B \cap P_\alpha v$ with $s_\alpha w > \phi(m(s_\alpha)v)$, a contradiction. Hence 2 holds for any representative $x \in v$.

Let then $x \in v$, let $l \in k \setminus \{0\}$ and let $x_1 = x_\alpha(l)xx_\alpha(-l) = u_1\dot{w}_1v_1$. Since $\alpha \in \Delta$ and $u, v \in P_\alpha^u$ we have $\dot{w}_1 = \dot{w}$ and $v_1 = (\dot{w}^{-1}x_\alpha(l)\dot{w})vx_\alpha(-l)$. The root α may not occur in v_1 . By Chevalley commutator formula this is possible only if $\dot{w}^{-1}x_\alpha(l)\dot{w} = x_\alpha(l)$, that is, only if 3 holds for X_α . Let $l \in k \setminus \{0\}$ and let $x_2 = x_{-\alpha}(l)xx_{-\alpha}(-l)$. Since $\alpha \in \Delta$ and $u, v \in P_\alpha^u$ we have $x_{-\alpha}(l)ux_{-\alpha}(-l) \in U$ so

$$x_2 \in U\dot{w}(\dot{w}^{-1}x_{-\alpha}(l)\dot{w})vx_{-\alpha}(-l).$$

If $(\dot{w}^{-1}x_{-\alpha}(l)\dot{w})vx_{-\alpha}(-l) \notin U$ we would have $\phi(B.x_2) \geq ws_\alpha \geq w$ hence $(\dot{w}^{-1}x_{-\alpha}(l)\dot{w})vx_{-\alpha}(-l) \in U$. By Chevalley commutator formula this is possible only if $\dot{w}^{-1}x_{-\alpha}(l)\dot{w} = x_{-\alpha}(l)$, that is, only if 3 holds for $X_{-\alpha}$.

Let x satisfy 1, 2, and 3. Then $P_\alpha v = P_\alpha x \subset B\dot{s}_\alpha X_\alpha.x \cup v$. Properties 2 and 3 imply that $B\dot{s}_\alpha X_\alpha.x \subset BwB$ and v is stationary. $\square$

Lemma 3.5 *Let $\mathcal{O}$ be a spherical conjugacy class, let $v \in \mathcal{V}$ be a stationary point and let $w = \phi(v)$. Let $\Pi = \{\alpha \in \Delta \mid w(\alpha) = \alpha\}$ and w_Π be the longest element in W_Π . Then $w = w_\Pi w_0$.*

Proof. By Theorem 2.8 the element $w \in W$ is an involution and by Lemma 3.4 if $w\alpha \in \Phi^+$ then $w\alpha = \alpha$ for every $\alpha \in \Delta$. The statement follows from [23, Proposition 3.5]. $\square$

Example 3.6 Let G be of type A_n and let $\mathcal{O}$ be a spherical conjugacy class. Then the image through ϕ of the dense B -orbit is $w = w_0 w_\Pi$ for some $\Pi \subset \Delta$. The set Π must be stabilized by $-w_0$ because for $\alpha \in \Pi$ we have

$$\alpha = w\alpha = w_0 w_\Pi \alpha \in -w_0 \Pi.$$

Besides, if α_j lies in Π then $\alpha_j = w(\alpha_j) = -w_\Pi(\alpha_{n-j+1})$ so α_j and α_{n-j} must lie in the same connected component of Π . Hence, $\Pi = \{\alpha_t, \alpha_{t+1}, \dots, \alpha_{n-t+1}\}$ for some t and

$$w = (s_{\beta_t} \cdots s_{\beta_{\lfloor \frac{n}{2} \rfloor}})w_0 = s_{\beta_1} \cdots s_{\beta_{t-1}}$$

where $\beta_1, \dots, \beta_{\lfloor \frac{n}{2} \rfloor}$ is the sequence given by the highest root, the highest root of the root system orthogonal to β_1 , and so further.

The example above shows that, for any stationary point, Π has to be invariant with respect to $-w_0$. Besides, the restriction of w_0 to $\Phi(\Pi)$ always coincides with w_Π .

If $w = \phi(v)$ for some stationary point $v \in \mathcal{V}$, the involution w may be written as a product of reflections with respect to mutually orthogonal roots $\gamma_1, \dots, \gamma_m$ so U_w (notation as in Section 1) is the subgroup generated by the root subgroups X_γ with $(\gamma, \gamma_j) = 0$ for every j . In other words, U_w is the subgroup generated by X_γ for $\gamma \in \Phi(\Pi)$ and U^w is normalized by U_w .

Lemma 3.7 *Let $\mathcal{O} = \mathcal{O}_x$ be a spherical conjugacy class in G . Let $v \in \mathcal{V}$ be a stationary point. If $x = u\dot{w}v \in U^w N(T)U$ then $B_x \subset uT^w U_w u^{-1}$.*

Proof. Let $g = v^w v_w t \in B_x$ with $v^w \in U^w$, $v_w \in U_w$ and $t \in T$, respectively. Then,

$$\begin{aligned} v^w v_w t u \dot{w} v &= v^w (v_w t u t^{-1} v_w^{-1}) \dot{w} (\dot{w}^{-1} v_w \dot{w}) (\dot{w}^{-1} t \dot{w}) v \\ &= u \dot{w} v v^w v_w t \in U^w \dot{w} B \end{aligned}$$

By normality of U^w and uniqueness of the decomposition we have

- $v^w = u(v_w t u^{-1} t^{-1} v_w^{-1})$ so v^w is completely determined by v_w , t and u ;
- $(\dot{w}^{-1} v_w \dot{w})(\dot{w}^{-1} t \dot{w}) v = v v^w v_w t$.

In particular $g = u v_w t u^{-1}$ hence $B_x \subseteq u U_w T u^{-1}$. Uniqueness of the decomposition in $B = TU$ implies that $\dot{w}^{-1} t \dot{w} = t$ so that $t \in T^w$ and $B_x \subseteq u T^w U_w u^{-1}$. $\square$

4 The dense B -orbit

We shall turn our attention to the special stationary point given by the dense B -orbit v_0 .

We will first analyze the possible Π for which $\phi(v_0) = w_0 w_\Pi$. We already know that they are subsets of Δ for which the restriction of w_0 to $\Phi(\Pi)$ coincides with the longest element of the parabolic subgroup W_Π of W . Next step will be to show which connected components of Π may not consist of isolated roots.

Lemma 4.1 *Let $\mathcal{O}$ be a spherical conjugacy class, let v_0 be its dense B -orbit and let $w = w_0 w_\Pi = \phi(v_0)$. There is no connected component in Π consisting of an isolated root $\alpha \in \Delta$ such that there is $\beta \in \Delta$ with the following properties:*

- β has the same length as α ;
- $w_0(\beta) = -\beta$;
- $\beta \not\perp \alpha$;
- $\beta \perp \alpha'$ for every $\alpha' \in \Pi \setminus \{\alpha\}$.

Proof. Let us choose a representative of v_0 of the form $x = \dot{w}v$ and let us assume that there are simple roots α and β with the above properties. The element w is the product of $w_0 s_\alpha$ with the longest element $w_{\Pi'}$ of the parabolic subgroup $W_{\Pi'}$ of W

associated with the complement Π' of α in Π . We claim that $v \in P_\beta^u$. Conversely, given a representative $\dot{s}_\beta$ of s_β in $N(T)$, we consider $y = \dot{s}_\beta \dot{w} v \dot{s}_\beta^{-1}$. Then, as s_β commutes with w_0 and $w_{\Pi'}$ by assumption, we would have, for some $l \in k \setminus \{0\}$:

$$y = \dot{w}_0 \dot{s}_{\alpha+\beta} \dot{w}_{\Pi'} v_1 x_{-\beta}(l) v_2 \in B w_0 s_{\alpha+\beta} w_{\Pi'} B s_\beta B; \quad v_i \in P_\beta^u.$$

Besides $w_0 s_{\alpha+\beta} w_{\Pi'}(\beta)$ is positive and different from β . Thus, $w_0 s_{\alpha+\beta} w_{\Pi'} s_\beta > w_0 s_{\alpha+\beta} w_{\Pi'}$ and in this case $\phi(B.y) = w_0 s_{\alpha+\beta} w_{\Pi'} s_\beta$ would not be an involution contradicting Lemma 2.8. Hence, $v \in P_\beta^u$ and $v' = \dot{s}_\beta v \dot{s}_\beta^{-1} \in U$.

Let us consider a representative $\dot{s}_\alpha$ of s_α in $N(T)$ and the element:

$$z = \dot{s}_\alpha \dot{s}_\beta \dot{w} v \dot{s}_\beta^{-1} \dot{s}_\alpha^{-1} = (\dot{w}_0 \dot{s}_\beta \dot{w}_{\Pi'}) (\dot{s}_\alpha \dot{s}_\beta v \dot{s}_\beta^{-1} \dot{s}_\alpha^{-1}) \in B w_0 s_\beta w_{\Pi'} B (\dot{s}_\alpha v' \dot{s}_\alpha^{-1}) B.$$

Here we have used that $w_0(\alpha) = -\alpha$ because α is an isolated root in Π . If $v' \in P_\alpha^u$ then $z \in B w_0 s_\beta w_{\Pi'} B$. If $v' \notin P_\alpha^u$ then

$$z \in B w_0 s_\beta w_{\Pi'} B (\dot{s}_\alpha v' \dot{s}_\alpha^{-1}) B \subset B w_0 s_\beta w_{\Pi'} B \cup B w_0 s_\beta s_\alpha w_{\Pi'} B.$$

It follows from Lemma 2.8 that also in this case $\phi(B.z) = w_0 s_\beta w_{\Pi'}$ because $(w_0 s_\beta s_\alpha w_{\Pi'})^2 = s_\alpha s_\beta \neq 1$. On the other hand, $\ell(\phi(B.z)) = \ell(w_0 s_\alpha w_{\Pi'}) = \phi(v_0)$ with $\phi(B.z) \neq \phi(v_0)$, contradicting $\phi(B.z) \leq \phi(v_0)$. $\square$

Corollary 4.2 *Let $\mathcal{O}$ be a noncentral spherical conjugacy class, let v_0 be its dense B -orbit and let $w = w_0 w_\Pi = \phi(v_0)$. Then Π is either empty or it is one of the following subsets of Δ :*

Type A_n

$$\circ \cdots \circ - - \bullet \cdots \bullet - - \circ \cdots \circ \quad \Pi = \{\alpha_l, \cdots, \alpha_{n-l+1}\}, \quad 2 \leq l \leq \left\lceil \frac{n}{2} \right\rceil$$

Type B_n

$$\circ \cdots \circ - - \bullet \cdots \bullet = > = \bullet \quad \Pi = \{\alpha_l, \cdots, \alpha_n\}, \quad 2 \leq l \leq n$$

$$\bullet - - \circ \cdots \bullet - - \circ - - \bullet \cdots \bullet = > = \bullet$$

$$\Pi = \{\alpha_1, \alpha_3, \cdots, \alpha_{2l-1}, \alpha_{2l+1}, \alpha_{2l+2}, \cdots, \alpha_n\}, \quad 1 \leq l \leq \frac{n}{2}$$

Type C_n

$$\circ \cdots \circ - - \bullet \cdots \bullet = < = \bullet \quad \Pi_1 = \{\alpha_l, \cdots, \alpha_n\}, \quad 2 \leq l \leq n$$

$$\bullet - - \circ \cdots \bullet - - \circ - - \bullet \cdots \bullet = < = \bullet$$

$$\Pi = \{\alpha_1, \alpha_3, \dots, \alpha_{2l-1}, \alpha_{2l+1}, \alpha_{2l+2}, \dots, \alpha_n\}, \quad 1 \leq l \leq \frac{n}{2}$$

Type D_n

$$\begin{array}{c} \circ \cdots \circ - - \bullet \cdots \bullet - - \bullet \\ | \\ \bullet \end{array} \quad \Pi_1 = \{\alpha_{2l+1}, \dots, \alpha_n\}, \quad 2 \leq l \leq \frac{n}{2} - 1$$

$$\begin{array}{c} \bullet - - \circ - - \bullet - - \circ \cdots \circ - - \bullet \cdots \bullet - - \bullet \\ | \\ \bullet \end{array}$$

$$\Pi = \{\alpha_1, \alpha_3, \dots, \alpha_{2l-1}, \alpha_{2l+1}, \alpha_{2l+2}, \dots, \alpha_n\}, \quad 1 \leq l < \frac{n}{2}$$

D_{2m}

$$\begin{array}{c} \bullet - - \circ - - \cdots - - \bullet - - \circ - - \bullet \\ | \\ \circ \end{array} \quad \Pi = \{\alpha_1, \alpha_3, \dots, \alpha_{2m-3}, \alpha_{2m-1}\}$$

$$\begin{array}{c} \bullet - - \circ - - \cdots - - \bullet - - \circ - - \circ \\ | \\ \bullet \end{array} \quad \Pi = \{\alpha_1, \alpha_3, \dots, \alpha_{2m-3}, \alpha_{2m}\}$$

D_{2m+1}

$$\begin{array}{c} \bullet - - \circ - - \cdots \bullet - - \circ - - \bullet - - \circ \\ | \\ \circ \end{array} \quad \Pi = \{\alpha_1, \alpha_3, \dots, \alpha_{2m-1}\}$$

Type E_6

$$\begin{array}{c} \bullet - - \bullet - - \bullet - - \bullet \\ | \\ \circ \end{array} \quad \Pi = \{\alpha_1, \alpha_3, \alpha_4, \alpha_5, \alpha_6\}$$

$$\begin{array}{c} \circ - - \bullet - - \bullet - - \bullet - - \circ \\ | \\ \circ \end{array} \quad \Pi = \{\alpha_3, \alpha_4, \alpha_5\}$$

Type E_7

$$\begin{array}{c} \circ - - \bullet - - \bullet - - \bullet - - \bullet \\ | \\ \bullet \end{array} \quad \Pi = \{\alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7\}$$

$$\begin{array}{c} \circ - - \bullet - - \bullet - - \bullet - - \circ - - \bullet \\ | \\ \bullet \end{array} \quad \Pi = \{\alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_7\}$$

$$\begin{array}{c} \circ - - \bullet - - \bullet - - \bullet - - \circ - - \circ \\ | \\ \bullet \end{array} \quad \Pi = \{\alpha_2, \alpha_3, \alpha_4, \alpha_5\}$$

$$\begin{array}{c} \circ - - \circ - - \circ - - \bullet - - \circ - - \bullet \\ | \\ \bullet \end{array} \quad \Pi = \{\alpha_2, \alpha_5, \alpha_7\}$$

Type E_8

$$\begin{array}{c} \bullet - - \bullet - - \bullet - - \bullet - - \bullet - - \bullet - - \circ \\ | \\ \bullet \end{array} \quad \Pi = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7\}$$

$$\begin{array}{c} \circ - - \bullet - - \bullet - - \bullet - - \bullet - - \circ \\ | \\ \bullet \end{array} \quad \Pi = \{\alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6, \alpha_7\}$$

$$\begin{array}{c} \circ - - \bullet - - \bullet - - \bullet - - \circ - - \circ - - \circ \\ | \\ \bullet \end{array} \quad \Pi = \{\alpha_2, \alpha_3, \alpha_4, \alpha_5\}$$

Type F_4

$$\bullet - - \bullet = > = \bullet - - \circ, \quad \Pi = \{\alpha_1, \alpha_2, \alpha_3\}$$

$$\circ - - \bullet = > = \bullet - - \bullet, \quad \Pi = \{\alpha_2, \alpha_3, \alpha_4\}$$

$$\circ - - \bullet = > = \bullet - - \circ, \quad \Pi = \{\alpha_2, \alpha_3\}$$

Type G_2

$$\bullet \equiv < \equiv \circ, \quad \Pi = \{\alpha_1\}$$

$$\circ \equiv < \equiv \bullet, \quad \Pi = \{\alpha_2\}.$$

Proof. Most of the restrictions are due to the fact that $w_\Pi = w_0|_{\Phi(\Pi)}$. The isolated roots occurring as connected components of Π are necessarily alternating, or differ by a node, or their length is different from the length of all adjacent roots, as $\{\alpha_n\}$ in type C_n . $\square$

Remark 4.3 All the above diagrams do actually occur when $k = \mathbb{C}$ (cfr. [8]) and they are strictly more than the Araki-Satake diagrams for symmetric conjugacy classes (see [1], [11], [20, Table 1]).

Theorem 4.4 Let $\mathcal{O}$ be a spherical conjugacy class, let v_0 be its dense B -orbit and let $w = \phi(v_0) = w_0 w_\Pi$. Then

$$\dim(\mathcal{O}) = \ell(w) + \text{rk}(1 - w).$$

Proof. We have divided the proof into a sequence of lemmas.

Let us recall that a standard parabolic subgroup can be naturally attached to $v \in \mathcal{V}$ ([13, §2]):

$$P(v) = \{g \in G \mid g.v = v\}.$$

Let $L(v)$ denote its Levi component containing T and let $\Delta(v)$ be the corresponding subset of Δ : this is the so-called set of simple roots of v . The derived subgroup $[L(v), L(v)]$ of $L(v)$ fixes a point in v (this is shown in [7, Lemma 1(ii)] and the argument works also in positive characteristic). We shall relate Π , $\Delta(v_0)$ and the centralizer of an element $x \in v_0$ in B .

Lemma 4.5 Let $\mathcal{O}$ be a spherical conjugacy class, let v_0 be its dense B -orbit and let $w = w_0 w_\Pi = \phi(v_0)$. Then $\Delta(v_0) \subset \Pi$.

Proof. If a simple root α occurs in $\Delta(v_0)$ then there is $x = uiv \in v$ for which $X_\alpha \in U \cap G_x$ and by Lemma 3.7 we have $1 \neq u^{-1} X_\alpha u \subset X_\alpha U^w \cap U_w$. This is possible only if $\alpha \in \Pi$ and u normalizes (hence centralizes) X_α . $\square$

Remark 4.6 In characteristic zero it follows from [7, Page 289] and [17, Corollary 3] that $\Pi = \Delta(v_0)$. We shall show that this is always the case.

Lemma 4.7 *Let $\mathcal{O}$ be a spherical conjugacy class, let v_0 be its dense B -orbit, let $x \in v_0$ and let $w = w_0 w_\Pi = \phi(v_0)$. If there exists $\alpha \in \Delta$ such that $X_\alpha \subset P_\alpha^u(P_\alpha)_x \cap U$ then $\alpha \in \Delta(v_0)$.*

Proof. If for every $r \in k$ there exists $u_r \in P_\alpha^u$ such that $g_r = x_\alpha(r)u_r \in (P_\alpha)_x$ this would hold for every choice of $x \in v_0$ so by Lemma 3.7 we have $\alpha \in \Pi$. Besides, the analysis of the B -orbits in $P_\alpha v_0$ shows that we are either in case I or in case IIa because the subgroup $H = \psi((P_\alpha)_x)$ would contain $\psi(X_\alpha)$. Since v_0 is the dense B -orbit, we cannot be in case IIa so $P_\alpha v_0 = v_0$ and $\alpha \in \Delta(v_0)$. $\square$

Let us recall that the depth $\text{dp}(\beta)$ of $\beta \in \Phi^+$ is the minimal length of a $\sigma \in W$ for which $\sigma(\beta) \in -\Phi^+$ ([4]). Then $\text{dp}(\beta) - 1$ is the minimal length of a $\sigma' \in W$ for which $\sigma'(\beta) \in \Delta$.

Lemma 4.8 *Let $\mathcal{O}$ be a spherical conjugacy class, let $v_0 \in \mathcal{V}$ be the dense B -orbit, let $\phi(v_0) = w_\Pi w_0$ and let $L(v_0)$ be the Levi factor of $P(v_0)$ containing T . There exists $x \in v_0$ such that*

1. x is centralized by $[L(v_0), L(v_0)]$;
2. x has decomposition $x = \dot{w}v \in U^w wB$;
3. no root in $\Phi^+(\Pi)$ occurs in the expression of v .

Proof. Let $x_1 \in v_0$ be an element that is centralized by $[L(v_0), L(v_0)]$. If $x_1 = u_1 \dot{w} v_1$ commutation with $X_{\pm\alpha}$ for $\alpha \in \Delta(v_0) \subset \Pi$ and Lemma 3.4 imply that $u_1, \dot{w}$ and v_1 are centralized by $X_{\pm\alpha}$, hence replacing x_1 by its B -conjugate $u_1^{-1} x u_1 = x = \dot{w} v$ yields the first and the second statement. Let us assume that for a fixed ordering of the positive roots the root $\gamma \in \Phi(\Pi)$ occurs in the expression of v and let us assume that γ is of minimal depth in $\Phi(\Pi)$ with this property. By Lemma 3.4 the root γ is not simple. Then, there exists $\sigma = s_{i_1} \cdots s_{i_r} \in W_\Pi$ such that $\sigma(\gamma) = \alpha \in \Pi$. Minimality of depth implies that for every other root $\gamma' \in \Phi(\Pi)$ occurring in v and every $j \geq 2$, we have $s_{i_j} s_{i_{j+1}} \cdots s_{i_r}(\gamma') \in \Phi^+$. Then for every representative $\dot{\sigma} \in N(T)$ we would have, for some $t \in T$,

$$x' = \dot{\sigma} x \dot{\sigma}^{-1} = \dot{w} t \dot{\sigma} v \dot{\sigma}^{-1} \in B w_\Pi w_0 B$$

so the B -orbit represented by x' would be stationary with $\dot{\sigma} v \dot{\sigma}^{-1} \notin P_\alpha^u$ contradicting Lemma 3.4. $\square$

Lemma 4.9 *Let $\mathcal{O}$ be a spherical conjugacy class, let v_0 be its dense B -orbit and let $w = w_0 w_\Pi = \phi(v_0)$. Then $\Delta(v_0)$ contains all long simple roots in Π . In particular, if Φ is simply laced $\Delta(v_0) = \Pi$ and there is $x \in v_0$ for which $U \cap G_x = U_w$.*

Proof. If $\Pi = \emptyset$ there is nothing to prove; if $\Pi = \Delta$ then $w = w_0 w_\Pi = 1 = \phi(v_0)$ so $\mathcal{O}$ is central, $P(v_0) = G$ and $\Pi = \Delta = \Delta(v_0)$. We shall assume throughout the proof that $\Pi \neq \emptyset, \Delta$.

Let $x = \dot{w}v$ be as in Lemma 4.8. If Π is strictly larger than $\Delta(v_0)$ by Lemma 4.7 then there exists $\alpha \in \Pi \setminus \Delta(v_0)$ such that X_α is not contained in the centralizer of x . Since $X_\alpha \subset G_{\dot{w}}$ by Lemma 3.4 we conclude that $X_\alpha \not\subset G_v$. By Chevalley's commutator formula this may happen only if there occurs in v a positive root γ with $\gamma + \alpha \in \Phi$. Indeed, in this case the structure constants $c_{\alpha, \gamma, 1, 1}$ in Chevalley's commutator formula ([21, Proposition 9.5.3]) would be nonzero. By Lemma 4.8 such a root γ does not lie in $\Phi^+(\Pi)$.

The basic idea of the proof is to show that if the above happens then there exists $v \in \mathcal{V}$ such that $\phi(v)$ is not an involution, contradicting Theorem 2.8. The proof consists in the construction of an element $\sigma = s_{i_t} \cdots s_{i_1} \in W$ such that:

1. $\sigma(\gamma) = -\alpha_{i_t}$;
2. the roots $\gamma'_j = s_{i_1} \cdots s_{i_j} \alpha_{i_{j+1}}$ for $j < t - 1$ do not occur in v ;
3. $\sigma w_0 w_\Pi \sigma^{-1}(\alpha_{i_t}) \in \Phi^+$ and is different from α_{i_t} .

The existence of a σ satisfying conditions 1 and 2 guarantees that for any representative $\dot{\sigma}$ of σ in $N(T)$ we have:

$$\dot{\sigma} x \dot{\sigma}^{-1} \in B \sigma w_0 w_\Pi \sigma^{-1} B X_{-\alpha_{i_t}} B = B \sigma w_0 w_\Pi \sigma^{-1} B s_{\alpha_{i_t}} B.$$

Condition 3 guarantees that $\sigma w_0 w_\Pi \sigma^{-1} < \sigma w w_\Pi \sigma^{-1} s_{i_t}$ and s_{i_t} does not commute with $\sigma w_0 w_\Pi \sigma^{-1}$. Then $\phi(B \dot{\sigma} x \dot{\sigma}^{-1}) = \sigma w_0 w_\Pi \sigma^{-1} s_{i_t}$ that is not an involution.

We will deal with the different possibilities for Π separately.

If Φ of type A_n then $\Pi = \{\alpha_l, \alpha_{l+1}, \dots, \alpha_{n-l+1}\}$ for $1 \leq l \leq \lfloor \frac{n+1}{2} \rfloor$. If $\alpha_i \notin \Delta(v_0)$ for some i with $l \leq i \leq n - l + 1$ then there occurs in v either $\mu_t = \sum_{j=t}^{i-1} \alpha_j$ with $t \leq l - 1$ or $\nu_t = \sum_{j=i+1}^t \alpha_j$ with $t \geq n - l + 2$. Let γ be a root of this form occurring in v and of minimal height. If $\gamma = \mu_t$ we consider $\sigma = \prod_{j=t}^{i-1} s_j$. Then $\sigma(\gamma) = -\alpha_t$ so condition 1 is satisfied. If for some r with $t < r \leq i - 1$ the root $\gamma_r = (\prod_{j=i-1}^{r-1} s_j) \alpha_r$ would occur in v we would contradict minimality of the height of γ so condition 2 is satisfied. Besides, if we put $\vartheta = -w_0$ we have:

$$\sigma w_0 w_\Pi \sigma^{-1}(\alpha_t) = \sigma w_\Pi \vartheta(\mu_t) = \alpha_{n-t+1} + \text{terms independent of } \alpha_{n-t+1}$$

so σ satisfies condition 3. The case $\gamma = \nu_t$ is treated similarly.

Let Φ be of type B_n . Then Π is either $\Pi_1 = \{\alpha_l, \dots, \alpha_n\}$ with $l \leq n$ of type B_{n-l+1} or A_1 ; or it is the union of Π_1 with $|\Pi_1|$ of the same parity as n and alternating isolated simple roots. The element w_Π is either w_{Π_1} or the product of the reflections corresponding to those isolated simple roots with w_{Π_1} .

Let us assume that there is $\alpha_i \in \Pi_1 \setminus \Delta(v_0)$ with $i \leq n-1$. Let $\gamma \in \Phi^+$ occur in v with $\alpha_i + \gamma \in \Phi$. It follows from Lemma 4.8 that γ is either $\mu_j = \sum_{p=j}^{i-1} \alpha_p$ for some $j < l$ or $\nu_j = \sum_{p=j}^i \alpha_p + 2 \sum_{q=i+1}^n \alpha_q$ for some $j < l$. Let us consider a μ_j with j maximal (that is, μ_j is of minimal height of this form). Then we can rule it out by using $\sigma_j = \prod_{p=j}^i s_p$. Condition 2 follows from minimality of the height of μ_j and Lemma 4.8, and condition 3 is checked observing that the coefficient of α_n in the expression of $\sigma_j w_\Pi(\gamma)$ is always positive. Let us then consider ν_j occurring in v with j maximal. Then we may rule it out by using $\tau_j = (\prod_{p=j}^n s_p)(\prod_{q=n-1}^{i+1} s_q)$. Condition 2 and 3 follow from maximality of j and Lemma 4.8 and from the fact that the coefficient of α_n in the expression of $\sigma_j w_\Pi(\gamma)$ is always positive. Thus $\Pi_1 \setminus \Delta(v_0) \subset \{\alpha_n\}$.

Let us now assume that Π is the union of Π_1 and alternating isolated simple roots. Suppose that α_i , for $1 \leq i < n$ is an isolated root in $\Pi \setminus \Delta(v_0)$. Then there occurs in v a γ that is either μ_j as above, with $j < i-1$; $\mu'_j = \sum_{p=i+1}^j \alpha_p$ for $i+1 < j \leq n$; ν_j as above, with $j < i$; or $\nu'_j = \sum_{p=i+1}^{j-1} \alpha_p + 2 \sum_{q=j}^n \alpha_q$ for $j \geq i+2$. The first set of roots is ruled out as in the case in which $\alpha_i \in \Pi_1$; the second set of roots is ruled out by using $\sigma'_j = \prod_{p=j}^{i+1} s_p$ and minimality of the height; the third set of roots is ruled out by τ_j as for $\alpha_i \in \Pi_1$. Here, for condition 2 the non-occurrence in v of the previous sets of roots is needed. Let ν'_j occur in v with j maximal. Then we consider $\tau'_j = (\prod_{q=j-1}^{n-1} s_q)(\prod_{p=n}^{i+1} s_p)$. Condition 2 is ensured by the maximality of j and by the non-occurrence of the roots previously excluded. Condition 3 holds because the coefficient of α_i in $\tau'_j w_\Pi(\nu'_j)$ is positive. Thus, all long roots in Π lie in $\Delta(v_0)$.

Let Φ be of type C_n . Then Π is $\Pi_1 = \{\alpha_l, \dots, \alpha_n\}$ with $l \leq n$ of type C_{n-l+1} or A_1 , or the union of Π_1 with alternating isolated simple short roots. If $\alpha_n \in \Pi_1 \setminus \Delta(v_0)$ then there would occur in v a root γ of the form $\mu_j = \sum_{p=j}^{n-1} \alpha_p$ with $j < l$. We rule this out by using $\sigma_j = \prod_{p=j}^{n-1} s_p$, hence $\alpha_n \in \Delta(v_0)$.

Let Φ be of type D_n . Then Π is either $\Pi_1 = \{\alpha_l, \dots, \alpha_{n-1}, \alpha_n\}$ for $1 \leq l \leq n-1$ with l odd; the union of Π_1 with alternating isolated simple roots; the set of all α_j with j odd; or the set of all α_j with j odd and $j \leq n-3$ or $j = n$.

If $\alpha_i \in \Pi_1 \setminus \Delta(v_0)$ for $i < n-1$ there would occur in v a root among the following: $\gamma_j = \sum_{p=j}^{i-1} \alpha_p$ for $j \leq l-1$; $\delta_j = \sum_{p=j}^i \alpha_p + 2 \sum_{q=i+1}^{n-2} \alpha_q + \alpha_{n-1} + \alpha_n$ for $j \leq l-1$. Let us consider γ_j of minimal height among the γ_t 's occurring in v . Then $\sigma_j = \prod_{p=j}^{i-1} s_p$ satisfies the necessary conditions for γ_j for

every choice of $\Pi \supset \Pi_1$ so there occurs no root of type γ_j in v . Let us consider the root of type δ_j of minimal height occurring in v . Then the Weyl group element $(\prod_{p=j}^n s_p)(\prod_{q=n-2}^{i+1} s_q)$ satisfies the necessary conditions in order to rule out δ_j . Therefore $\alpha_i \in \Delta(v_0)$ for every $l \leq i \leq n-2$. If $\alpha_{n-1} \in \Pi \setminus \Delta(v_0)$ there would occur in v a root among the following: $\gamma_j = \sum_{p=j}^{n-2} \alpha_p$ for $j \leq l-1$; or $\delta'_j = \sum_{p=j}^{n-2} \alpha_p + \alpha_n$ for $j < l$. The roots of type γ_j are ruled out as for α_i for $i < n-1$. In order to rule out the roots of type δ'_j we consider the minimal occurring in v and we apply $\tau_j = (\prod_{p=j}^{n-2} s_p)s_n$. Condition 3 is ensured when $\Pi_1 \subset \Pi$ because $j < l$ and when Π is a set of alternating roots because the coefficient of α_{n-2} in $(\prod_{p=j}^{n-2} s_p)s_n w_\Pi(\delta'_j)$ is always positive. Thus, $\alpha_{n-1} \in \Delta(v_0)$ and $\alpha_n \in \Delta(v_0)$ by symmetry.

Let α_i be an isolated root in $\Pi \setminus \Delta(v_0)$ for $i \leq n-2$. Then there would occur in v a root of type γ_j for $j < i$, $\mu_j = \sum_{p=i+1}^j \alpha_p$ for $i+1 \leq j \leq n-1$, $\mu_n = \sum_{p=i+1}^{n-2} \alpha_p + \alpha_n$, $\mu'_n = s_n \mu_{n-1}$, $\nu_j = \sum_{p=i+1}^{j-1} \alpha_p + 2 \sum_{q=j}^{n-2} \alpha_q + \alpha_{n-1} + \alpha_n$ for $j \geq i+2$, or δ_j for $j < i$. These roots are handled making use, in this order, of: σ_j for γ_j ; $\omega_j = \prod_{p=j}^{i+1} s_p$ for μ_j for $j \leq n-1$; $\omega_n = s_n(\prod_{p=n-2}^{i+1} s_p)$ for μ_n ; $\omega'_n = s_n \omega_{n-1}$ for μ'_n ; $(\prod_{p=j}^n s_p)(\prod_{q=n-2}^{i+1} s_q)$ for ν_j ; τ_j for δ_j .

In all cases the required conditions are easily verified. In particular, condition 3 holds, for every choice of Π for which α_i is an isolated root, because either α_i , α_{i-1} or α_{i+1} will occur in $\sigma w_\Pi \gamma$ with positive coefficient, for σ any of the above Weyl group elements. This concludes the proof for Φ of type D_n .

Let Φ be of type E_6 , so that Π is either $\Pi_1 = \{\alpha_1, \alpha_3, \alpha_4, \alpha_5, \alpha_6\}$ or $\Pi_2 = \{\alpha_3, \alpha_4, \alpha_5\}$. For both cases the analysis in type A_n shows that if $\Pi \neq \Delta(v_0)$ then the roots γ occurring in v for which $\gamma + \alpha \in \Phi$ for some $\alpha \in \Pi$ may not lie in the root subsystem of type A_5 generated by Π_1 .

Let $\Pi = \Pi_1$ and let us assume that $\alpha_6 \notin \Pi$. Then there occurs γ in v where γ is one of the following roots: $\mu_1 = \alpha_2 + \alpha_4 + \alpha_5$, $\mu_2 = s_3 \mu_1$, $\mu_3 = s_4 \mu_2$, $\mu_4 = s_1 \mu_2$, $\mu_5 = s_4 \mu_4$, $\mu_6 = s_3 \mu_5$. All these roots can be ruled out by using, respectively: $\sigma_1 = s_2 s_4 s_5$, $\sigma_2 = \sigma_1 s_3$, $\sigma_3 = \sigma_2 s_4$, $\sigma_4 = \sigma_2 s_1$, $\sigma_5 = \sigma_4 s_4$, $\sigma_6 = \sigma_5 s_3$. Condition 3 is always satisfied because α_6 occurs in the expression of $\sigma_i w_\Pi(\mu_i)$ with positive coefficient for every i . Therefore, α_6 (and, symmetrically, α_1) lie in $\Delta(v_0)$ and all roots occurring in v are orthogonal to them because $[L(v_0), L(v_0)]$ centralizes x .

Let us assume that $\alpha_4 \notin \Delta(v_0)$. The roots α_2 and $\alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4 + 2\alpha_5 + \alpha_6$ are the only positive roots that do not lie in $\Phi(\Pi)$, are orthogonal to α_1 and α_6 and are such that $\gamma + \alpha_4 \in \Phi$. The first can be ruled out with $\sigma = s_2$ while for the second we may use $\tau = s_2 s_4 s_3 s_5 s_1 s_3 s_6 s_5$. Since there are no positive roots γ orthogonal to α_1 , α_4 and α_6 , that do not lie in $\Phi(\Pi)$ and for which $\gamma + \alpha_3$ or $\gamma + \alpha_5 \in \Phi$ we conclude that $\Pi = \Delta(v_0)$ and we have the statement for $\Pi = \Pi_1$.

Let $\Pi = \Pi_2$ and let us assume that $\alpha_3 \notin \Delta(v_0)$. Then there would occur in v a root γ among the following roots: $\nu_1 = \alpha_2 + \alpha_4$, $\nu_2 = s_5\nu_1$, $\nu_3 = s_6\nu_2$, $\nu_4 = \sum_{j=1}^5 \alpha_j + \alpha_4$, $\nu_5 = s_6\nu_4$, $\nu_6 = s_5\nu_5$. They can be ruled out by using $\sigma_1 = s_2s_4$, $\sigma_2 = \sigma_1s_5$, $\sigma_3 = \sigma_2s_6$, $\sigma_4 = \sigma_2s_3s_4s_1$, $\sigma_5 = \sigma_4s_6$ and $\sigma_6 = \sigma_5s_5$, respectively. Here condition 3 follows because the coefficient of α_5 in $\sigma_i w_\Pi \nu_i$ is positive. Thus α_3 and, by symmetry, α_5 lie in $\Delta(v_0)$ and all roots occurring in v are orthogonal to them. If α_4 would not lie in $\Delta(v_0)$ there would occur in v either α_2 or $\sum_{j=1}^6 \alpha_j$. The first root is ruled out through s_2 while the second is ruled out through $s_2s_3s_4s_5s_1s_6$ and we have the statement in type E_6 .

Let Φ be of type E_7 . Then Π corresponds either to a subdiagram Π_1 of type D_6 (containing all simple roots but α_1), a subdiagram Π_2 of type D_4 (containing $\alpha_2, \alpha_3, \alpha_4, \alpha_5$), the union of Π_2 and α_7 , or $\Pi_3 = \{\alpha_2, \alpha_5, \alpha_7\}$.

Let $\Pi = \Pi_1$ and let us assume that $\alpha_3 \notin \Delta(v_0)$. Then there would occur in v one of the following roots: α_1 ; $\gamma_1 = \alpha_4 + \sum_{i=1}^5 \alpha_i$; $\gamma_2 = s_6\gamma_1$; $\gamma_3 = s_5\gamma_2$; $\gamma_4 = s_7\gamma_2$; $\gamma_5 = s_5\gamma_4$; $\gamma_6 = s_6\gamma_5$; $\gamma_7 = \alpha_1 + 2\alpha_2 + 2\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7$. None of these roots may actually occur, as it can be shown through s_1 ; $\sigma_1 = s_1s_3s_4s_2s_5s_4$; $\sigma_2 = \sigma_1s_6$; $\sigma_3 = \sigma_2s_5$; $\sigma_4 = \sigma_2s_7$; $\sigma_5 = \sigma_4s_5$; $\sigma_6 = \sigma_5s_6$ and $\sigma_7 = \sigma_1s_3(\prod_{j=6}^4 s_j)s_2(\prod_{p=7}^4 s_p)$, respectively. Condition 2 holds because all simple reflections except from the first one in the decomposition of σ_j lie in W_Π . Condition 3 holds for all σ_j because the coefficient of α_7 is positive in $\sigma_j w_\Pi \gamma_j$. Therefore $\alpha_3 \in \Delta(v_0)$ and all roots in v are orthogonal to it. If α_2 would not lie in $\Delta(v_0)$ there would occur in v one of the following roots: $\delta_1 = \alpha_1 + \alpha_3 + \alpha_4$; $\delta_2 = s_5\delta_1$; $\delta_3 = s_6\delta_2$; $\delta_4 = s_7\delta_3$; $\delta_5 = \alpha_1 + \alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6$; $\delta_6 = s_7\delta_5$; $\delta_7 = s_6\delta_6$; $\delta_8 = s_5\delta_7$. None of them may occur, as it is shown through $\tau_1 = s_1s_3s_4$; $\tau_2 = \tau_1s_5$; $\tau_3 = \tau_2s_6$; $\tau_4 = \tau_3s_7$; $\tau_5 = \tau_1s_2(\prod_{j=5}^3 s_j)(\prod_{p=6}^4 s_p)$. It follows that $\alpha_2 \in \Delta(v_0)$. Since the only positive root that is orthogonal to α_2 and α_3 and lies in $\Phi \setminus \Phi(\Pi_1)$ is the highest root in $\Phi(E_7)$, we have the statement for Π_1 .

Let Π be either Π_2 or the union of Π_2 and α_7 . The discussion concerning type D_6 shows that the sum of a root in $\Phi(\Pi_1)$ occurring in v with a root in Π_2 is never a root. If $\alpha_3 \notin \Delta(v_0)$ there would occur in v some of the γ_i 's that have been discussed for $\Pi = \Pi_1$. The Weyl group elements s_i ; σ_i for $i = 1, 2$ still satisfy the necessary conditions. In order to rule out γ_3 we may use $s_1s_3\sigma_3$; for γ_4 we may use $\sigma'_4 = (\prod_{j=4}^3 s_j)(\prod_{p=5}^4 s_p)(\prod_{j=6}^4 s_j)(\prod_{p=7}^5 s_p)s_2s_1s_4$; for γ_5 we may use $\sigma'_5 = s_3s_1s_4s_2(\prod_{j=5}^4 s_j)(\prod_{p=6}^5 s_p)(\prod_{j=7}^6 s_j)s_4s_5$; for γ_6 we may use $\sigma'_6 = \sigma'_5s_5s_4$; for γ_7 we may use $\sigma'_7 = \sigma_1s_3s_1(\prod_{j=6}^4 s_j)s_2(\prod_{p=7}^4 s_p)$. The necessary conditions hold for both choices of Π . Thus, $\alpha_3 \in \Delta(v_0)$. If $\alpha_2 \notin \Delta(v_0)$ there would occur in v one of the δ_j 's we mentioned when discussing $\Pi = \Pi_1$. We may use τ_i for $i = 1, 2, 3$ to exclude the first three roots;

for τ_4 condition 2 is no longer satisfied and we use $\tau'_4 = (\prod_{j=7}^3 s_j)s_1$; for δ_j for $j = 5, 6$ we may use $\tau'_j = s_2s_4s_3s_1(\prod_{p=j}^3 s_p)(\prod_{q=j+1}^4 s_q)$; for δ_7 we use $\tau'_7 = s_2s_4s_3s_1s_5s_4(\prod_{p=6}^3 s_p)(\prod_{q=7}^4 s_q)$. Therefore, $\alpha_2 \in \Delta(v_0)$. Arguing as we did for $\Pi = \Pi_1$ we conclude that all X_α for $\alpha \in \Pi$ commute with $\dot{w}v$.

Let $\Pi = \{\alpha_2, \alpha_5, \alpha_7\}$. It follows from the discussion concerning D_6 that all roots γ occurring in v and such that $\gamma + \alpha \in \Phi$ for some $\alpha \in \Pi$ may not lie in the root subsystem $\Phi(\Pi_1)$. Let us assume that $\alpha_7 \notin \Delta(v_0)$. Then there would occur in v one of the following roots: $\mu_1 = \alpha_1 + \sum_{j=3}^6 \alpha_j$, $\mu_2 = s_2\mu_1$, $\mu_3 = s_4\mu_2$, $\mu_4 = s_3\mu_3$, $\mu_5 = s_5\mu_3$, $\mu_6 = s_3\mu_5$, $\mu_7 = s_4\mu_6$, $\mu_8 = s_2\mu_7$. None of these roots may occur since we may use, respectively: $\sigma_1 = s_1 \prod_{i=3}^6 s_i$; $\sigma_2 = \sigma_1 s_2$; $\sigma_3 = s_1(\prod_{i=3}^5 s_i)s_2(\prod_{i=4}^6 s_i)$; $\sigma_4 = (\prod_{i=1}^5 s_i)s_4(\prod_{i=2}^4 s_i)s_6$; $\sigma_5 = (\prod_{i=1}^5 s_i)s_4s_6s_5$; $\sigma_6 = (\prod_{i=5}^2 s_i)s_4s_5s_1(\prod_{i=3}^6 s_i)$; $\sigma_7 = (\prod_{i=1}^6 s_i)s_4s_3s_5s_4$; and $\sigma_8 = (\prod_{i=5}^6 (s_i s_{i-1}))s_3s_4(\prod_{i=1}^6 s_i)s_3s_4s_2$. Then $\alpha_7 \in \Delta(v_0)$ so all roots occurring in v are orthogonal to it. Let us assume that $\alpha_2 \notin \Delta(v_0)$. The possible positive roots ν that are orthogonal to α_7 , do not lie in $\Phi(\Pi_1)$ and for which $\alpha_2 + \nu \in \Phi$ are: $\nu_1 = \alpha_1 + \alpha_3 + \alpha_4$, $\nu_2 = s_5\nu_1$, $\nu_3 = \alpha_1 + \alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + 2\alpha_6 + \alpha_7$, $\nu_4 = s_5\nu_3$. These roots are excluded by using $\tau_1 = s_1s_3s_4$, $\tau_2 = \tau_1s_5$, $\tau_3 = (\prod_{i=4}^1 s_i)(\prod_{i=5}^7 s_i s_{i-1})s_3s_4$ and $\tau_4 = s_3s_1s_4s_2s_5s_4s_3(\prod_{i=6}^7 s_i s_{i-1})s_4s_5$. Thus $\alpha_2 \in \Delta(v_0)$ and all roots occurring in v are orthogonal to α_2 and α_7 . Since there are no positive roots γ in $\Phi \setminus \Phi(\Pi_1)$ such that $\gamma \perp \alpha_2, \alpha_7$ and $\gamma + \alpha_5 \in \Phi$ we conclude that $\alpha_5 \in \Delta(v_0)$ and we have the statement for E_7 .

Let Φ be of type E_8 . Then Π corresponds either to a subdiagram Π_0 of type E_7 (all simple roots but α_8), to a subdiagram Π_1 of type D_6 (all simple roots but α_1, α_8), a subdiagram Π_2 of type D_4 (consisting of $\alpha_2, \alpha_3, \alpha_4, \alpha_5$).

If $\Pi = \Pi_0$ and $\alpha_3 \notin \Pi_0$ there would occur in v one of the following roots: $\gamma_1 = \sum_{j=4}^8 \alpha_j$; $\gamma_2 = s_2\gamma_1$; $\gamma_3 = 2\alpha_1 + 2\alpha_2 + 3\alpha_3 + 5\alpha_4 + 4\alpha_5 + 3\alpha_6 + 2\alpha_7 + \alpha_8$; $\gamma_4 = s_2\gamma_3$; $\mu_1 = \alpha_4 + \sum_{j=1}^8 \alpha_j$; $\mu_2 = s_5\mu_1$; $\mu_3 = s_6\mu_2$; $\mu_4 = s_7\mu_3$; $\mu_5 = \alpha_1 + 2\alpha_2 + 2\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7 + \alpha_8$, $\mu_6 = s_7\mu_5$; $\mu_7 = s_6\mu_6$; $\mu_8 = s_5\mu_7$.

We exclude γ_1 and γ_2 through $\sigma_1 = \prod_{j=8}^4 s_j$ and $\sigma_2 = \sigma_1 s_2$; γ_3 through $\sigma_3 = (\prod_{p=8}^1 s_p)s_4s_5s_3s_4s_2(\prod_{p=6}^3 s_p)(\prod_{j=7}^4 s_j)s_1$ and γ_4 through $\sigma_4 = \sigma_3 s_2$. We rule out μ_j for $1 \leq j \leq 4$ through $\tau_1 = \sigma_2 s_3 s_1 s_4$; $\tau_2 = \tau_1 s_5$; $\tau_3 = \tau_2 s_6$; $\tau_4 = \tau_3 s_7$. We exclude μ_j for $j = 5, 6, 7, 8$ through $\tau_5 = (\prod_{p=7}^1 s_p)s_8 s_7 s_4 s_3 s_5 s_4 s_2(\prod_{q=6}^4 s_q)$, $\tau_6 = \tau_5 s_7$, $\tau_7 = \tau_6 s_6$ and $\tau_8 = \tau_7 s_5$. In order to verify condition 2 we use Lemma 4.8 and the non-occurrence of the previously excluded roots. Therefore $\alpha_3 \in \Delta(v_0)$ and all roots occurring in v are orthogonal to it. If $\alpha_2 \notin \Delta(v_0)$ there would occur in v one of the following roots: $\beta_1 = \alpha_1 + \sum_{j=3}^8 \alpha_j$; $\beta_2 = \alpha_1 + \alpha_2 + 2\alpha_3 + 3\alpha_4 + 2\alpha_5 + \alpha_6 + \alpha_7 + \alpha_8$; $\beta_3 = s_6\beta_2$; $\beta_4 = s_7\beta_3$; $\beta_5 = s_5\beta_4$; $\beta_6 = s_6\beta_5$; $\beta_7 = s_5\beta_3$; $\beta_8 = \alpha_1 + 2\alpha_2 + 3\alpha_3 + 5\alpha_4 + 4\alpha_5 + 3\alpha_6 + 2\alpha_7 + \alpha_8$. All these roots may be ruled out by using, respectively, $\omega_1 = (\prod_{j=8}^3 s_j)s_1$; $\omega_2 =$

$\omega_1 s_2 s_4 s_3 s_5 s_4$; $\omega_3 = \omega_2 s_6$; $\omega_4 = \omega_3 s_7$; $\omega_5 = \omega_4 s_5$; $\omega_6 = \omega_5 s_6$; $\omega_7 = \omega_3 s_5$; $\omega_8 = s_3 s_1 (\prod_{p=4}^6 s_p s_{p-1}) s_2 s_4 s_3 s_1 s_7 s_8 (\prod_{q=5}^4 s_q s_{q+1} s_{q+2}) s_2 (\prod_{q=4}^5 s_p s_{p-1})$.

Thus, α_2 and α_3 both lie in $\Delta(v_0)$ and all roots occurring in v are orthogonal to them. If $\alpha_5 \notin \Delta(v_0)$ there would occur in v one of the following roots: $\nu_1 = \sum_{j=6}^8 \alpha_j$; $\nu_2 = \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \nu_1$; $\nu_3 = 2\alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 3\alpha_5 + 3\alpha_6 + 2\alpha_7 + \alpha_8$; $\nu_4 = \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \nu_3$. They are ruled out with $r_1 = \prod_{j=8}^6 s_j$; $r_2 = (\prod_{j=8}^2 s_j) s_4$; $r_3 = (\prod_{j=8}^1 s_j) s_4 s_3 s_5 s_4 s_2 (\prod_{p=6}^3 s_p) (s_1 s_7 s_6)$; $r_4 = r_3 (s_5 s_4 s_3 s_2 s_4)$. Thus all roots in v are orthogonal to α_5 as well. The possible roots γ occurring in v such that $\alpha_6 + \gamma \in \Phi$ are then $\alpha_8 + \alpha_7$, that is ruled out by $s_8 s_7$, and $2\alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + 2\alpha_7 + \alpha_8$, that is ruled out by $\sigma = (\prod_{j=8}^1 s_j) s_4 s_3 s_5 s_4 s_2 s_6 s_7 (\prod_{p=5}^3 s_p) s_1$. The roots γ that might occur in v for which $\alpha_7 + \gamma \in \Phi$ are α_8 , that is ruled out by s_8 , and $2\alpha_1 + 2\alpha_2 + 3\alpha_3 + 4\alpha_4 + 3\alpha_5 + 2\alpha_6 + \alpha_7 + \alpha_8$ which is ruled out by σs_7 . Thus $\alpha_j \in \Delta(v_0)$ for $j = 2, 3, 5, 6, 7$. The only positive root that does not lie in $\Phi(E_7)$ and that is orthogonal to these simple roots is the highest root in Φ , whence the statement.

Let us assume that $\Pi = \Pi_1$. As a consequence of the discussion concerning type E_7 if $\gamma + \alpha \in \Phi$ with $\alpha \in \Pi$ then γ may not lie in the root subsystem corresponding to Π_0 . Let us assume that $\alpha_3 \notin \Pi$ so that a root among the γ_j , $1 \leq j \leq 4$ and μ_p for $1 \leq p \leq 8$ introduced when dealing with $\Pi = \Pi_0$ would occur in v . The Weyl group elements σ_1 and σ_2 exclude γ_1 and γ_2 . In order to exclude γ_3 and γ_4 we use, respectively, $\sigma'_3 = s_8 s_7 \sigma_3$ and $\sigma'_4 = \sigma'_3 s_2$. Here, in order to verify condition 2 we use the non-occurrence of γ_1, γ_2 . The roots μ_j for $1 \leq j \leq 8$ are excluded by the same Weyl group elements as for $\Pi = \Pi_0$, therefore $\alpha_3 \in \Delta(v_0)$ and all roots occurring in v are orthogonal to it. If $\alpha_2 \notin \Delta(v_0)$ there would occur in v one of the β_j 's that were introduced in the discussion of $\Pi = \Pi_0$. The roots β_j for $1 \leq j \leq 8$ are excluded by the same Weyl group elements as for $\Pi = \Pi_0$. If $\alpha_5 \notin \Delta(v_0)$ there would occur in v one of the ν_i for $i = 1, 2, 3, 4$. The first two roots are excluded by the same Weyl group elements r_1 and r_2 as in case $\Pi = \Pi_0$, where we use for condition 2 that all roots occurring in v are orthogonal to α_2 and α_3 . In order to exclude ν_3 and ν_4 we use $r'_3 = s_8 s_7 r_3$ and $r'_4 = s_8 s_7 r_4$, respectively. If $\alpha_6 \notin \Delta(v_0)$ then the possible roots occurring in v that added to α_6 provide a root are ruled out by $s_8 s_7$, and $\sigma' = s_8 s_7 \sigma$, with notation as for $\Pi = \Pi_0$. Then the roots γ for which $\alpha_7 + \gamma \in \Phi$ are those indicated for $\Pi = \Pi_0$ and are ruled out by s_8 and σs_7 . As there are no roots γ in $\Phi \setminus \Phi(\Pi_0)$ that are orthogonal to $\alpha_2, \alpha_3, \alpha_5, \alpha_6$ and for which $\alpha_4 + \gamma \in \Phi$ we have the statement for $\Pi = \Pi_1$.

Let $\Pi = \Pi_2$. It follows from the discussion concerning Φ of type E_7 that if γ occurs in v and $\gamma + \alpha \in \Phi$ for some $\alpha \in \Pi$ then γ may not lie in the root subsystem corresponding to Π_0 . If $\alpha_3 \notin \Delta(v_0)$ there would occur in v a root among the γ_j 's and the μ_j 's that we have previously discussed. Then γ_1 and γ_2 are excluded by σ_1 and σ_2 ; γ_3 is excluded by σ'_3 . We rule out μ_j for $j = 1, 2, 3$

with the same Weyl group elements τ_j that we used for $\Pi = \Pi_0$. Then μ_4 is excluded by $\tau'_4 = (\prod_{p=7}^3 s_p s_{p+1}) s_2 (\prod_{q=4}^7 s_p s_{p-1}) s_1$ and we can rule out γ_4 by $s_1 (\prod_{p=3}^8 s_p) s_2 (\prod_{q=4}^7 s_q) (\prod_{p=3}^6 s_p) s_2 s_4 s_5 s_3 s_4 s_2 s_1$. The roots μ_j , for $5 \leq j \leq 8$ are ruled out, respectively, by:

$$\tau'_5 = (\prod_{p=4}^8 s_p s_{p-1}) s_2 s_4 s_3 s_1 s_5 s_6 s_3 s_4 s_5 s_2 s_4;$$

$$\tau'_6 = (\prod_{p=4}^8 s_p s_{p-1}) s_2 s_4 s_3 s_1 s_6 s_5 s_7 s_6 s_3 s_4 s_5 s_2 s_4;$$

$\tau'_7 = \tau'_6 s_6$ and $\tau'_8 = s_1 s_3 s_4 s_2 (\prod_{p=5}^3 s_p) s_1 (\prod_{q=6}^8 s_q) \prod_{p=5}^4 (s_p s_{p+1}) s_2 s_4 \prod_{q=7}^5 s_q$. Thus $\alpha_3 \in \Delta(v_0)$ and all roots occurring in v are orthogonal to it. If $\alpha_2 \notin \Delta(v_0)$ there would occur in v one of the β_j 's previously introduced. We exclude the β_j for $1 \leq j \leq 3$ and $j = 7$ by the same Weyl group elements ω_j as for $\Pi = \Pi_0$ and $\Pi = \Pi_1$. Then we may exclude β_j for $j = 4, 5, 6$ making use of $\omega'_4 = (\prod_{p=3}^8 s_p) s_2 s_4 s_3 s_1 (\prod_{q=5}^7 s_q s_{q-1}) s_3 s_4$; $\omega'_5 = \omega'_4 s_5$ and $\omega'_6 = \omega'_5 s_6$. Finally, we exclude β_8 with the same Weyl group element ω_8 that we used when $\Pi = \Pi_0$. The order in which we exclude roots allows the control on condition 2. Thus, α_2 and $\alpha_3 \in \Delta(v_0)$ and if $\gamma \in \Phi$ occurs in v and $\gamma + \alpha_5 \in \Phi$ then $\gamma = \nu_i$ for some i , with notation as for $\Pi = \Pi_0$. The first three roots are ruled out by the Weyl group elements r_1, r_2, r'_3 that we introduced in the previous cases. The root ν_4 is ruled out by $r''_4 = s_2 s_4 s_3 s_5 s_4 s_1 s_3 (\prod_{p=6}^2 s_p) s_4 s_7 s_8 (\prod_{q=6}^3 s_q) s_1 (\prod_{p=7}^2 s_p) s_4$. Thus, all roots occurring in v are orthogonal to $\alpha_5, \alpha_2, \alpha_3$. No root γ in $\Phi \setminus \Phi(\Pi_0)$ for which $\alpha_4 + \gamma \in \Phi$ satisfies these conditions, whence the statement in type E_8 .

Let Φ be of type F_4 . Then Π is either of type B_3, C_3 , or B_2 . It follows from the discussion in types B_n and C_n that all roots γ occurring in v and for which $\alpha + \gamma \in \Phi$ for some long root $\alpha \in \Pi$ may not lie in any root subsystem of type C_3 or B_3 containing Π .

If $\Pi = \{\alpha_1, \alpha_2, \alpha_3\}$ is of type B_3 and $\alpha_1 \notin \Delta(v_0)$ there would occur in v either $\nu_1 = \alpha_2 + \alpha_3 + \alpha_4$, $\nu_2 = \alpha_3 + \nu_1$, or $\nu_3 = \alpha_4 + \nu_2$. We rule them out with $\sigma_1 = s_4 s_3 s_2$, $\sigma_2 = \sigma_1 s_3$, $\sigma_3 = s_2 s_3 s_2 \sigma_1$ so $\alpha_1 \in \Delta(v_0)$ and all roots occurring in v are orthogonal to it. If $\alpha_2 \notin \Delta(v_0)$, there occurs in v either $\alpha_3 + \alpha_4$ or $\alpha_1 + 2\alpha_2 + 4\alpha_3 + 2\alpha_2$. We rule them out through $s_4 s_3$ and $s_2 s_3 s_4 s_1 s_1 s_2 s_3$, respectively. Hence, all long roots in Π lie in $\Delta(v_0)$.

If $\Pi = \{\alpha_2, \alpha_3, \alpha_4\}$ and $\alpha_2 \notin \Delta(v_0)$ there would occur in v one of the following roots: $\alpha_1, \gamma_1 = \alpha_1 + \alpha_2 + 2\alpha_3, \gamma_2 = \gamma_1 + \alpha_4, \gamma_3 = s_4 \gamma_1, \gamma_4 = \alpha_1 + 2\alpha_2 + 4\alpha_3 + 2\alpha_4$. These roots are excluded by $s_1, \sigma_1 = s_1 s_2 s_3, \sigma_2 = s_3 s_2 s_1 s_2 s_4 s_3, \sigma_3 = \sigma_1 s_4, \sigma_4 = s_2 s_3 s_1 s_2 s_4 s_3$ so $\alpha_2 \in \Delta(v_0)$.

If $\Pi = \{\alpha_2, \alpha_3\}$, then it is contained in a subsystem of type C_3 and in a subsystem of type B_3 and we may make use of the discussion of type B_n and C_n . If $\alpha_2 \notin \Delta(v_0)$ there would occur in v either $\alpha_1 + \alpha_2 + 2\alpha_3 + \alpha_4$ or $\alpha_1 + 2\alpha_2 + 4\alpha_3 + 2\alpha_4$. The first root is excluded by $\sigma = s_3 s_2 s_1 s_2 s_4 s_3$ and the second is excluded by $s_2 \sigma$.

Let Φ be of type G_2 . Then Π consists of a simple root. If $\Pi = \{\alpha_1\}$ there is nothing to prove. If $\Pi = \{\alpha_2\}$ and $\Delta(v_0) = \emptyset$ there occurs in v a root γ with $\gamma \in \{\alpha_1, 3\alpha_1 + \alpha_2\}$. We may then use $\sigma = s_1; s_2s_1$, respectively, to rule out these possibilities.

The last statement follows from the fact that there is $x \in v_0$ that is centralized by $[L(\Pi), L(\Pi)]$, so that $U \cap G_x$ is the unipotent subgroup generated by the root subgroups corresponding to roots in Π . $\square$

We have shown so far that $\Pi \setminus \Delta(v_0)$ consists at least of all long roots in Π . The method used in the proof of Lemma 4.9 might fail for those short roots $\alpha \in \Pi$ for which the α -string through some positive root γ is of length at least 3. When its length is 3, that is, in the doubly-laced case, the root γ might be orthogonal to Π so condition 3 might fail. In type G_2 condition 2 might fail for all $\sigma \in W$. Next Lemma will deal with these cases replacing the representative $x \in v_0$ if needed.

Lemma 4.10 *Let Φ be of type B_n, C_n, F_4 or G_2 , let $\mathcal{O}$ be a spherical conjugacy class, let v_0 be its dense B -orbit and let $w = w_0w_\Pi = \phi(v_0)$. Then $\Delta(v_0) = \Pi$ and there is $x \in v_0$ for which $G_x \cap U = U_w$.*

Proof. Let $x = \dot{w}v \in v_0$ and let α be a short root in Π . We will show that no positive root ν with $\nu + \alpha \in \Phi$ may occur in v . For those roots ν with $\nu - \alpha \notin \Phi$ we will argue as in Lemma 4.9. For the other roots we will show that $\nu + \alpha$ may not occur in any X_α -conjugate of x . Then Chevalley commutator formula implies that ν may not occur in v .

We shall deal with the different root systems separately and we will use the terminology and notation introduced in Lemma 4.9.

Let Φ be of type B_n . The only short root in Π is $\alpha = \alpha_n$. Let us assume that $\alpha_n \notin \Delta(v_0)$. The root subgroups that do not commute with X_{α_n} correspond to the roots $\gamma_j = \sum_{p=j}^{n-1} \alpha_p$ and $\delta_j = \gamma_j + \alpha_n$ for $j < l$. The roots γ_j are ruled out by using $\sigma_j = \prod_{p=j}^{n-1} s_p$ and the usual minimality argument. If Π is the union of Π_1 and alternating isolated long roots then for every j there exists $\alpha \in \Delta(v_0)$ that is not orthogonal to δ_j , so none of them may occur in v . Therefore if $\alpha_n \notin \Delta(v_0)$ we necessarily have $\Pi = \Pi_1$ and some root δ_j occurs in v . Let $a \in k$ and let us consider $x_1 = x_{\alpha_n}(a)xx_{\alpha_n}(-a) = \dot{w}v_1$. We claim that $\delta_j + \alpha_n$ may not occur in v_1 for any choice of $a \in k$ and for any j . Indeed, if $j \geq l$ then the argument in the proof of Lemma 4.8 shows that $\delta_j + \alpha_n \in \Phi(\Pi)$ may not occur in v_1 . Let $\delta_j + \alpha_n$ be of minimal height occurring in v_1 . Then $\delta_j + \alpha_n$ is ruled out by using $\sigma_j s_n$. If some δ_j would occur in v then there would exist $a \in k$ for which $\delta_j + \alpha_n$ would occur in $x_{\alpha_n}(a)v x_{\alpha_n}(-a) = v_1$, leading to a contradiction. The statement is proved for Φ of type B_n .

Let Φ be of type C_n and let $\Pi_1 = \{\alpha_l, \dots, \alpha_n\}$ with $l \leq n$. If $\alpha_i \in \Pi_1 \setminus \Delta(v_0)$ then $i < n$ and there occurs γ in v among the roots: $\mu_{ji} = \sum_{p=j}^{i-1} \alpha_p$ with $j < l$ and $\nu_{ji} = \sum_{p=j}^i \alpha_p + 2 \sum_{q=i+1}^{n-1} \alpha_q + \alpha_n$ for $j < l$. The first set of roots is ruled through $\sigma_{ji} = \prod_{p=j}^{i-1} s_p$ while the second set of roots is ruled out through $\tau_{ji} = (\prod_{p=j}^{n-1} s_p)(\prod_{q=n}^{i+1} s_q)$. Thus, $\Pi_1 \subset \Delta(v_0)$.

Let Π be the union of Π_1 and alternating isolated simple roots and let us assume that one of these, say α_i , does not lie in $\Delta(v_0)$. Then there occurs in v one of the roots: μ_{ji} as before, for $j < l$; $\mu'_{ji} = \sum_{p=i+1}^j \alpha_p$ for $j \geq i+2$; $\delta_{ji} = \sum_{p=i+1}^{j-1} \alpha_p + 2 \sum_{q=i}^{n-1} \alpha_q + \alpha_n$ for $j \geq i+2$; ν_{ji} as before for $j \leq i$; $\nu_{ii} - \alpha_i$.

The roots of type μ_{ji} are ruled out as when $\alpha_i \in \Pi_1$; the roots of type μ'_{ji} are ruled out by using $\sigma'_{ji} = \prod_{p=j}^{i+1} s_p$; the roots of type δ_{ji} are ruled out through $\omega_{ji} = (\prod_{p=j-1}^{n-1} s_p)(\prod_{q=n}^{i+1} s_q)$ where, for condition 2, we use the non-occurrence of any μ'_{ji} in v . Then, the roots of type ν_{ji} for $j < i$ are ruled out as when $\alpha_i \in \Pi_1$. Here, in order to verify condition 2 we use the non-occurrence in v of the previous sets of roots. We see that the root $\nu_{ii} - \alpha_i$ may not occur in v by using $\prod_{p=n}^{i+1} s_p$. Hence, $\alpha_i \notin \Delta(v_0)$ if and only if ν_{ii} occurs in v . Let $a \in k$ and let $x_1 = x_{\alpha_i}(a) \dot{w} x_{\alpha_i}(-a) = \dot{w} v_1$. We claim that $\nu_{jj} + \alpha_j$ may not occur in v_1 for any $a \in k$ and for any j . Indeed, if $j \geq l$ we may rule it out as in Lemma 4.8. If $j < l$ we consider $\nu_{jj} + \alpha_j$ of minimal height and we rule it out by using $\prod_{p=n}^j s_p$. This implies that ν_{ii} may not occur in v so the statement holds for Φ of type C_n .

Let Φ be of type F_4 and let us assume that $\Pi = \{\alpha_1, \alpha_2, \alpha_3\}$. By Lemma 4.9 the roots occurring in v are orthogonal to α_1 and α_2 . If $\alpha_3 \notin \Delta(v_0)$ there would occur in v one of the following roots: α_4 and $\nu = \alpha_1 + 2\alpha_2 + 3\alpha_3 + 2\alpha_4$. The first root is ruled out by s_4 . If ν occurred in v , there would be an X_{α_3} -conjugate $x_1 = \dot{w} v_1$ of x for which $\nu + \alpha_3$ occurs in v_1 . This is ruled out by using $\sigma = s_2 s_3 s_4 s_1 s_2 s_3$. Condition 2 is ensured by Lemma 4.8 and by Lemma 4.9, where we showed that $\alpha_2 + \alpha_3 + \alpha_4$ and $\alpha_2 + 2\alpha_3 + \alpha_4$ do not occur in v . Condition 1 and 3 are straightforward. Hence $\Delta(v_0) = \{\alpha_1, \alpha_2, \alpha_3\}$.

Let us assume that $\Pi = \{\alpha_2, \alpha_3\}$. The roots occurring in v have to be orthogonal to α_2 . If $\alpha_3 \notin \Delta(v_0)$ there would occur in v one of the following roots: α_4 , $\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4$; $\mu = \alpha_1 + \alpha_2 + \alpha_3$ or ν as in the previous case. The first and the second root are ruled out, respectively, by s_4 and $s_3 s_4 s_2 s_1$. Then $\mu + \alpha_3$ and $\nu + \alpha_3$ may not occur in any X_{α_3} -conjugate of v because they are ruled out, respectively, by $s_1 s_2 s_3$ and σ , whence $\Delta(v_0) = \Pi$.

Let us assume that $\Pi = \{\alpha_2, \alpha_3, \alpha_4\}$ and that $\alpha_3 \notin \Delta(v_0)$. The roots occurring in v have to be orthogonal to α_2 and may not lie in $\Phi(\Pi)$. Therefore there would occur in v one of the following roots: $\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4$; μ or ν as in the previous case. The first root is ruled out by $s_3 s_2 s_1 s_4$. The remaining two roots are handled

as in the previous case: $\mu + \alpha_3$ and $\nu + \alpha_3$ may not occur in any X_{α_3} -conjugate of v because they are ruled out by the same Weyl group element as for $\Pi = \{\alpha_2, \alpha_3\}$. Thus, $\alpha_3 \in \Delta(v_0)$ and all roots occurring in v are orthogonal to it. There are no roots γ that are orthogonal to α_3, α_2 , do not lie in Π and are such that $\gamma + \alpha_4 \in \Phi$, whence the statement for Φ of type F_4 .

Let Φ be of type G_2 let $\Pi = \{\alpha_1\}$. If $\Delta(v_0) = \emptyset$ there would occur in v a root $\gamma \in \{\alpha_2, \alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2\}$. The first two roots are ruled out through $\sigma = s_2; s_1 s_2$, respectively. Hence $\alpha_1 \notin \Pi$ if and only if $2\alpha_1 + \alpha_2$ occurs in v . On the other hand, $3\alpha_1 + \alpha_2$ may not occur in any X_{α_1} -conjugate of v because it is ruled out by $s_2 s_1$. This concludes the proof for Φ of type G_2 . $\square$

Lemma 4.11 *Let G be a simple algebraic group and suppose that the longest element $w_0 \in W$ acts as -1 in its standard representation. Let $\mathcal{O}$ be a spherical conjugacy class, let v_0 be its dense B -orbit and let $w = w_0 w_\Pi = \phi(v_0)$. Then there is $x \in v_0$ for which $U \cap G_x = U_w$ and $T_x^\circ = (T^w)^\circ$ so that $\dim T_x = \text{rk}(1 - w)$.*

Proof. Let $\Gamma(\mathcal{O})$ be the free abelian subgroup of all characters of B that arise as weights of eigenvectors of B in the function field $k(\mathcal{O})$. Let $r(\mathcal{O})$ denote its rank as a free abelian group. By [17, Corollary 1, Corollary 2] in characteristic 0 or [14, Lemma 2.1] for arbitrary characteristic and by Lemmas 4.9, 4.10 we have $\dim(\mathcal{O}) = \ell(w) + r(\mathcal{O})$. Since, for a representative $x \in v_0$ we have $\dim(B_x) \leq \dim T^w + \dim U_w$ we have

$$(4.1) \quad r(\mathcal{O}) \geq \text{rk}(1 - w)$$

where the rank of $1 - w$ is the minimal number of reflections needed in the decomposition of w (cfr. [18, Page 910]), that is, the dimension of the eigenspace corresponding to -1 .

Besides, by [7, Lemma 1(iii)] the set Π is orthogonal to $\Gamma(\mathcal{O})$, whence

$$(4.2) \quad r(\mathcal{O}) \leq n - |\Pi|.$$

If $w_0 = -1$ the restriction of w_Π to $\Phi(\Pi)$ is also -1 . Then the -1 eigenspace of w_Π has dimension at least $|\Pi|$ so the dimension of the -1 eigenspace of $w = w_0 w_\Pi$ is at most $n - |\Pi|$. This is indeed the exact dimension, for if the dimension of the 1 -eigenspace of w would be greater than $|\Pi|$, there would be χ in $\mathbb{Q}\Delta \setminus \mathbb{Q}\Pi$ such that

$$\chi = \sum_{\alpha \in \Pi} c_\alpha \alpha + \sum_{\beta \in \Delta \setminus \Pi} d_\beta \beta = w_0 w_\Pi(\chi) = \sum_{\alpha \in \Pi} c_\alpha \alpha - \sum_{\beta \in \Delta \setminus \Pi} d_\beta w_\Pi(\beta).$$

We may write w_Π as a product of reflections with respect to orthogonal roots belonging to $\Phi(\Pi)$ so that $w_\Pi(\beta) \in \beta + \mathbb{Q}(\Pi)$. If $w_0 w_\Pi(\chi) = \chi$ we would have

$2 \sum_{\beta \in \Delta \setminus \Pi} d_\beta \beta \in \mathbb{Q}\Pi$, a contradiction. Thus, $\text{rk}(1 - w) = n - |\Pi|$. Combining this with (4.1) and (4.2) we obtain the statement. $\square$

Lemma 4.12 *Let G be a simple algebraic group with Dynkin diagram of type A_n , D_{2m+1} , or E_6 . Let $\mathcal{O}$ be a spherical conjugacy class, let v_0 be its dense B -orbit and let $w = w_0 w_\Pi = \phi(v_0)$. Then there is $x \in v_0$ for which $U \cap G_x = U_w$ and $T_x^\circ = (T^w)^\circ$ so that $\dim T.x = \text{rk}(1 - w)$.*

Proof. Let $x = v$ be as in Lemma 4.8. Since it is centralized by $[L(v_0), L(v_0)]$ we see that $\alpha^\vee(h) \in T_x$ for every $\alpha \in \Pi$. Let ϑ be the non-trivial automorphism of Φ that is equal to $-w_0$. Then $(T^w)^\circ$ is generated by $\alpha^\vee(h)$ for $\alpha \in \Pi$ and $\alpha^\vee(h)(\vartheta\alpha)^\vee(h^{-1})$ for $\alpha \in \Delta \setminus \Pi$. If $(T^w)^\circ \not\subset T_x$ there would be a γ occurring in v with $(\gamma, \alpha) \neq (\gamma, \vartheta\alpha)$ for some $\alpha \in \Delta \setminus \Pi$. We will use the same argument as in the proof of Lemma 4.9 to show that this cannot be the case.

Let Φ be of type A_n . Then $\Pi = \{\alpha_l, \dots, \alpha_{n-l+1}\}$ and by Lemma 4.9 we have $\gamma \perp \Pi$. It follows that $\gamma = \alpha_j + \dots + \alpha_t$ with $t \neq n - j + 1$ and either $j \leq l - 1$ and $t \geq n - l + 1$; or $j < t \leq l - 2$; or $n - l + 3 \leq j < t$. Let γ be the root of minimal height of the first type. If $t > n - j + 1$ we consider $\sigma = s_j s_{j+1} \dots s_t$. Condition 1 is satisfied by construction. If $s_p \dots s_t \gamma' = \alpha_{p-1}$ for $p - 1 > j$ then $\gamma' = \alpha_{p-1} + \dots + \alpha_t$ with $n - t + 1 < j < p$ so minimality of the height of γ forces condition 2 to hold. Condition 3 is satisfied because α_{n-t+1} appears in the expression of $\sigma w_0 w_\Pi(\gamma)$ with positive coefficient. If $t < n - j + 1$ we shall use σ^{-1} . If $j < t \leq l - 2$ or $n - l + 3 \leq j < t$ the same argument for γ minimal applies with $\sigma = s_j s_{j+1} \dots s_t$ and the conditions are easier to verify. This argument holds also if $\Pi = \emptyset$.

Let Φ be of type D_{2m+1} . If $\Pi_1 \neq \emptyset$ (notation as in Lemma 4.9) then any positive root with $(\gamma, \alpha) \neq (\gamma, \vartheta\alpha)$ is of the form $\gamma = \sum_{p=j}^{n-2} \alpha_p + \alpha_q$ for $q = 2m, 2m + 1$ and none of these roots is orthogonal to $\Pi_1 \subset \Pi$.

If $\Pi_1 = \emptyset$ we consider γ of minimal height among the roots γ described above occurring in v and $\sigma = s_j \dots s_{n-2} s_q$. Then conditions 1 and 2 are satisfied by construction and minimality of height and condition 3 follows because α_q occurs with positive coefficient in $\sigma w_0 w_\Pi \gamma$.

Let Φ be of type E_6 . If γ is a positive root for which $(\gamma, \alpha_1) \neq (\gamma, \alpha_6)$ then either γ or $\vartheta(\gamma)$ lies in $\{\alpha_1, \alpha_3, \alpha_1 + \alpha_3, \alpha_3 + \alpha_4, \alpha_2 + \alpha_3 + \alpha_4, \alpha_1 + \alpha_3 + \alpha_4, \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4, \alpha_2 + \alpha_3 + \alpha_4, \alpha_2 + \alpha_3 + 2\alpha_4 + 2\alpha_5 + \alpha_6, \alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + 2\alpha_5 + \alpha_6\}$. We shall rule out these roots, their image through ϑ can be handled symmetrically. The roots α_i with $i = 1, 3$ may not occur either because $\alpha_i \in \Pi$, or because $\alpha_i \notin \Pi$, or because they are ruled out using s_i . Besides, $\gamma \perp \Pi$ hence $\vartheta\gamma \perp \Pi$ because Π is ϑ -stable. We may rule out the remaining roots by using $\sigma = s_3 s_1, s_4 s_3, s_2 s_4 s_3, s_4 s_3 s_1, s_2 s_4 s_3 s_1, s_3 s_4 s_2 s_5 s_4 s_6 s_5, s_3 s_4 s_2 s_5 s_4 s_6 s_5 s_1$. The conditions in the proof

of Lemma 4.9 are easily verified. Let us observe that some roots need only be handled when $\Pi = \emptyset$, that is, when $w = w_0$. Thus if a positive root γ is such that $(\gamma, \alpha) \neq (\gamma, \vartheta\alpha)$ for some $\alpha \in \Delta$ we necessarily have $\alpha = \alpha_3$ or $\alpha = \vartheta(\alpha_3)$ and either γ or $\vartheta(\gamma)$ lies in $\{\gamma_1 = \alpha_1 + \alpha_3 + \alpha_4 + \alpha_5, \gamma_2 = \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 + \alpha_5, \gamma_3 = \alpha_2 + \alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6\}$. These roots can only occur for $\Pi = \emptyset$ and in this case we may use, respectively, $\sigma_1 = s_5s_4s_3s_1$; $\sigma_2 = s_2s_4s_3s_1s_5$ and $\sigma_3 = s_5s_4s_2s_3s_4s_5s_6$. By symmetry we deduce that no root with $\gamma \neq \vartheta\gamma$ may occur in $\mathfrak{v}$. This concludes the proof of the Lemma. $\square$

It follows from Lemmas 4.9, 4.10, 4.11 and 4.12 that there is $x \in v_0$ such that $(B_x)^\circ = (T^w)^\circ U_w$ and this concludes the proof of Theorem 4.4. $\square$

Corollary 4.13 *Let $\mathcal{O}$ be a spherical conjugacy class, let $v_0 \in \mathcal{V}$ be the dense B -orbit and let $w = w_0w_\Pi = \phi(v_0)$. Then $v_0 = \mathcal{O} \cap BwB$.*

Proof. The arguments in Lemmas 4.8 depend only on the image of a B -orbit through ϕ . If for some $v \in \mathcal{V}$ we have $\phi(v) = w$ there is a representative $x = \dot{w}v$ of v that is centralized by $[L(v), L(v)]$ with no occurrence of roots in $\Phi(\Pi)$ in $\mathfrak{v}$. The centralizer of x in U will be contained in U_w and $\Delta(v) \subset \Pi$. Proceeding as in Lemmas 4.9 and 4.10 we conclude that $\Pi = \Delta(v) = \Delta(v_0)$. The arguments in the proof of Lemmas 4.11 and 4.12 involve only the image of ϕ , hence $\dim v = \dim v_0$, whence $v = v_0$. $\square$

Remark 4.14 If $\mathcal{O}$ is a symmetric conjugacy class over an algebraically closed field of odd or zero characteristic Theorem 4.4 follows from [19, Proposition 3.9, Theorem 4.6, Theorem 7.1] and Corollary 4.13 follows from [19, Theorem 7.11, Lemma 7.12, Theorem 7.13]. If $\mathcal{O}$ is a spherical conjugacy class over an algebraically closed field of characteristic zero Theorem 4.4 is [8, Theorem 1] and Corollary 4.13 is [8, Corollary 26].

Corollary 4.15 *Let $\mathcal{O}$ be a spherical conjugacy class with dense B -orbit v_0 and $\phi(v_0) = w_0w_\Pi$. Then there is $x = \dot{w}v \in v_0$ for which all roots occurring in $\mathfrak{v}$ are fixed by ϑ and orthogonal to Π .*

Corollary 4.16 *Let λ be a weight of an irreducible representation occurring in the G -module decomposition of the ring of regular functions on $\mathcal{O}$ over $\mathbb{C}$. Then $\lambda^* = -w_0\lambda = \lambda$. In particular, $\Gamma(\mathcal{O})$ is symmetric.*

Proof. By [17, §6] we have $\Gamma(\mathcal{O}) = \text{Ann}(T_x)$. It follows from Corollary 4.15 and Lemmas 4.11, 4.12 that $(\lambda, \alpha^\vee) = (\lambda, \vartheta\alpha^\vee)$ for $\alpha \in \Delta \setminus \Pi$ and $0 = (\lambda, \alpha^\vee) = (\lambda, -\vartheta\alpha^\vee) = (\lambda, \vartheta\alpha^\vee)$ for $\alpha \in \Pi$. Hence $\vartheta\lambda = -w_0\lambda = \lambda$. $\square$

Corollary 4.17 *Let λ be a weight of an irreducible representation occurring in the G -module decomposition of the ring of regular functions on $\mathcal{O}$ over $\mathbb{C}$. Let v_0 be the dense B -orbit and $w = w_0 w_\Pi = \phi(v_0)$. Then $\lambda \in \text{Ker}(1 + w) \cap P^+ \cap Q$.*

Proof. The inclusion in P^+ is obvious, the inclusion in Q follows from the fact that the ring of regular functions on $\mathcal{O}$ is a G_{ad} -module. It follows from Corollary 4.17 that $-\lambda = w_0 \lambda = w w_\Pi \lambda = w \lambda$ since $\lambda \perp \Pi$. $\square$

References

- [1] S. ARAKI, *On root systems and an infinitesimal classification of irreducible symmetric spaces*, J. Math. Osaka City Univ. 13 , 1–34 (1962).
- [2] A. BOREL, *Linear Algebraic Groups* W. A. Benjamin, Inc. (1969).
- [3] N. BOURBAKI, *Éléments de Mathématique. Groupes et Algèbres de Lie, Chapitres 4,5, et 6*, Masson, Paris (1981).
- [4] A. BJÖRNER, F. BRENTI, *Combinatorics of Coxeter Groups*, Springer (2005).
- [5] M. BRION, *Quelques propriétés des espaces homogènes sphériques*, Manuscripta Math. 55, 191–198 (1986).
- [6] M. BRION, *The behaviour at infinity of the Bruhat decomposition*, Comment. Math. Helv. 73, 137–174 (1998).
- [7] M. BRION, *On orbit closures of spherical subgroups in flag varieties*, Comment. Math. Helv. 76, 263–299 (2001).
- [8] N. CANTARINI, G. CARNOVALE, M. COSTANTINI, *Spherical orbits and representations of $\mathcal{U}_\varepsilon(\mathfrak{g})$* Transformation Groups, 10, No. 1, 29–62 (2005).
- [9] C. DE CONCINI, V. G. KAC, C. PROCESI, *Quantum coadjoint action*, Journal of the Amer. Math. Soc. 5, 151–190 (1992).
- [10] F. GROSSHANS, *Contractions of the actions of reductive algebraic groups in arbitrary characteristic*, Invent. Math. 107, 127–133 (1992).
- [11] A. G. HELMINCK *Algebraic groups with a commuting pair of involutions and semisimple symmetric spaces*, Adv. Math. 71, 21–91 (1988).
- [12] J. HUMPHREYS *Reflection Groups and Coxeter Groups*, Cambridge University Press (1990).

- [13] F. KNOP, *The asymptotic behavior of invariant collective motion*, Invent. Math. 116, 309–328 (1994).
- [14] F. KNOP, *On the set of orbits for a Borel subgroup*, Comment. Math. Helvetici 70, 285–309 (1995).
- [15] J.G.M. MARS, T. A. SPRINGER, *Hecke algebras representations related to spherical varieties*, Representation Theory, 2, 33–69 (1998).
- [16] A. MELNIKOV, *Description of B-orbit closures of order 2 in upper-triangular matrices*, Transform. Groups, 11(2), 217–247 (2006).
- [17] D. PANYUSHEV *Complexity and rank of homogeneous spaces*, Geom. Dedicata, 34, 249–269 (1990).
- [18] S. PASIENCIER, H-C. WANG *Commutators in a semi-simple Lie group*, Proc. Amer. Math. Soc. 13(6), 907–913 (1962).
- [19] R. W. RICHARDSON, T.A. SPRINGER, *The Bruhat order on symmetric varieties*, Geom. Dedicata, 35(1-3), 389–436 (1990).
- [20] T.A. SPRINGER *The classification of involutions of simple algebraic groups* J. Fac. Sci. Univ. Tokyo, Sect. IA, Math. 34, 655–670, (1987).
- [21] T.A. SPRINGER *Linear Algebraic Groups, Second Edition* Progress in Mathematics 9, Birkhäuser (1998).
- [22] T.A. SPRINGER, *Schubert varieties and generalizations*, Representation theories and algebraic geometry (Montreal, PQ, 1997), 413–440, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 514, Kluwer Acad. Publ., Dordrecht, (1998).
- [23] T.A. SPRINGER, *Some results on algebraic groups with involutions*, Algebraic groups and related topics (Kyoto/Nagoya, 1983), 525–543, Adv. Stud. Pure Math., 6, North-Holland, Amsterdam, (1985).
- [24] T.A. SPRINGER, R. STEINBERG *Conjugacy classes*, In: “Seminar on algebraic groups and related finite groups”. LNM 131, 167–266, Springer-Verlag, Berlin Heidelberg New York (1970).
- [25] E. VINBERG, *Complexity of action of reductive groups*, Func. Anal. Appl. 20, 1–11 (1986).